

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1972

[Full Table of Content](#)



### 72/R/WW/12 Winter Wheat Effects of Blue Green Algae

#### Rothamsted Research

Rothamsted Research (1973) *72/R/WW/12 Winter Wheat Effects of Blue Green Algae* ; Yields Of The Field Experiments 1972, pp 290 - 292 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/WW/12

WINTER WHEAT

EFFECTS OF BLUE-GREEN ALGAE

Object: To study the establishment of blue-green algae in relation to culture type, method of application, irrigation and nitrogen fertiliser. Effects on yield of wheat are also tested - Great Field II.

Sponsor: P.J. Dart.

Design: A single replicate of 5 x 3 x 2 x 2 in 60 plots.

Whole plot dimensions: 2.13 x 4.57. Area harvested: 0.00081.

Treatments: All combinations of:-

1. Type of algal culture\* (C):
  - None (0)
  - Nostoc elliposporium (isolated from Broadbalk Field) (1)
  - Nostoc punctiformum (isolated from Broadbalk Field) (2)
  - Anabaena cylindrica (3)
  - A mixture of the three species (4)
2. Irrigation (I): None (0), 50 mm (2 inches) during the first week in July (1), soil surface kept continually moist (2).
3. Method of application of algal inoculum\* (M): In water as a spray (1), as a dry sand crust broadcast by hand (2).
4. N levels: 0, 50 kg as 'Nitro-Chalk'.

\* Rates of application of inoculum were as follows:

	(in g/ha of dry weight of algae)	
Spray inoculum:	C1:	920
	C2:	920
	C3:	420
	C4:	670
Dry sand inoculum: (estimated weights)	C1:	450
	C2:	830
	C3:	260
	C4:	830

Basal applications: 280 kg (0:20:20) combine drilled.

Seed: Maris Ranger sown at 200 kg.

72/R/WW/12

Cultivations, etc.: Ploughed: 23 Sept, 1971. Seed combine drilled:  
6 Oct. N and water cultures applied: 9 May, 1972. Sand cultures  
applied: 15 May. Combine harvested: 31 Aug. Previous crops:  
Barley 1970 beans 1971.

NOTE: Estimates were made of the extent of algal cover and species  
involved and of recovery of inoculum types. N fixation was  
measured by acetylene reduction assay.

Standard error per plot (estimated from 4 factor interaction).  
Grain, tonnes/hectare: 0.543 or 8.2% (8 d.f.)

72/R/WW/12

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	I			M		N: KG/HA		Mean
	0	1	2	1	2	0	50	
C								
0	6.23	6.60	6.92	6.70	6.47	5.70	7.47	6.58
1	7.07	6.48	6.76	6.75	6.78	6.00	7.54	6.77
2	6.47	6.37	6.44	6.67	6.18	5.38	7.47	6.42
3	6.74	6.59	6.73	6.72	6.65	5.64	7.74	6.69
4	6.77	6.46	6.45	6.61	6.52	5.76	7.36	6.56
		I						
		0		6.77	6.54	5.80	7.51	6.66
		1		6.60	6.40	5.54	7.46	6.50
		2		6.69	6.63	5.75	7.57	6.66
				M				
				1		5.79	7.59	6.69
				2		5.60	7.44	6.52
Mean						5.70	7.51	6.61

STANDARD ERRORS OF DIFFERENCES

C	I	M	N	C	C	I	C	I	M
				I	M	M	N	N	N
0.222	0.172	0.140	0.140	0.384	0.314	0.243	0.314	0.243	0.198

Mean D.M. %: 85.8