Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1972



Full Table of Content

# 72/R/WS/4 Spring Wheat Dwarf Varieties, N and Ccc

#### **Rothamsted Research**

Rothamsted Research (1973) 72/R/WS/4 Spring Wheat Dwarf Varieties, N and Ccc; Yields Of The Field Experiments 1972, pp 297 - 298 - DOI: https://doi.org/10.23637/ERADOC-1-62

#### 72/R/WS/4

#### SPRING WHEAT

#### DWARF VARIETIES, N AND CCC

Object: To study the effects of CCC (chlormequat) and a range of nitrogen levels on four semi-dwarf spring wheat varieties - Fosters Corner.

Sponsor: S.A.W. French.

Design: A single replicate of 4 x 4 x 4 in 4 blocks of 4 whole plots split into 4 sub plots, with split plot confounding of 3 factor interaction with blocks.

Whole plot dimensions: 2.16 x 29.6. Sub plot area harvested: 0.00080.

Treatments: All combinations of:Whole plots: 1. Varieties: Charles Peguy\* (CP), Tobari (TO),
6WUK3 (W3), 6WUK4(W4).

\* Formerly Benoist 257.

Sub plots: 2. N levels: 75, 150, 225, 300 kg as 'Nitro-Chalk'.

3. CCC: None, 1.1, 2.2, 3.3 kg in 340 l.

Basal applications: 630 kg (0:14:28). Weedkiller: Bromoxynil, ioxynil, dichlorprop and MCPA ('Tetroxone' at 5.6 l in 340 l).

Seed: Sown at 190 kg.

Cultivations, etc.: Deep-time cultivated twice: 12 Nov, 1971.

Basal PK applied: 21 Mar, 1972. Seed drilled: 23 Mar. Weedkiller applied: 17 May. N applied: 1 May. CCC applied: 1 June.

Combine harvested: 7 Sept. Previous crops: Fallow 1970, potatoes 1971.

NOTE: Plant counts were made shortly after germination, shoot counts in early June, and ear counts at ear emergence. Crop height was measured at harvest.

Standard errors per plot.

Grain, tonnes/hectare: Whole plot: 0.193 or 4.4% (6 d.f.)

Sub plot: 0.276 or 6.3% (18 d.f.)

# 72/R/WS/4

#### TABLES OF MEANS

## GRAIN: TONNES/HECTARE

N: KG/HA

CCC: KG/HA

	75	150	225	300	0.0	1.1	2.2	3.3	Mean
VARIETY		1	onle	100			- deda	24 (5)	2,8 -0
CP TO W3 W4	5.61 4.58 2.33 4.12	5.47 4.89 2.41 4.75	5.12 4.93 2.72 4.92	4.93 4.96 2.66 5.11	4.85	5.19 4.57 2.51 4.82	5.04 5.06 2.70 4.84	5.37 4.88 2.58 4.51	5.28 4.84 2.53 4.72
	455	N	: KG/E	IA	n duen			No.	i ecis
SECONDA	ge) - Ya	150 4.41 4.33 4.54 4.23 225 4.53 4.17 4.44 4.55	4.23 4.23 4.55 4.34	4.16 4.38 4.42 4.42					
Mean	Eftergrein	MD 04	oper 6	est, in	4.36	4.27	4.41	4.34	4.34

Mean D.M. %: 84.2

### STANDARD ERRORS OF DIFFERENCES

	VARIETY	N	CCC	VARIETY N	VARIETY	CCC
Unless same	0.137	0.097	0.097	0.217	0.217	0.195
VARIETY				0.195	0.195	