

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1972

[Full Table of Content](#)



### 72/R/B/1 and 72/R/BB/B/1 Barley Growth and Yield on Contrasted Sites

#### Rothamsted Research

Rothamsted Research (1973) *72/R/B/1 and 72/R/BB/B/1 Barley Growth and Yield on Contrasted Sites* ; Yields Of The Field Experiments 1972, pp 299 - 302 - DOI:  
<https://doi.org/10.23637/ERADOC-1-62>

72/R/B/1 and 72/BB/B/1

BARLEY

GROWTH AND YIELD ON CONTRASTED SITES

**Object:** To try to account for yields and differences between yields of barley on sites at Rothamsted and Broom's Barn by studying crop growth rates, nutrient uptake, water use etc., at a wide range of nitrogen levels, with and without irrigation. Also to study the interaction between site differences and crops (see also 72/R & BB/WW/6) - Rothamsted (R), Pastures and Broom's Barn (BB), Windbreak Field.

**Sponsors:** P.J. Welbank, F.V. Widdowson.

**Design:** 3 randomised blocks of 2 plots, split into 6.

**Whole plot dimensions:**

Pastures (R): 15.2 x 48.0. Sub plot area harvested: 0.00434.

Windbreak Field (BB): 36.6 x 45.7. Sub plot area harvested: 0.00413.

**Treatments:** All combinations of:-

Whole plots: 1. Irrigation (I): None (0), full irrigation (C).

Sub plots: 2. Nitrogen: 31, 63, 94, 125, 157, 188 kg N as 'Nitro-Chalk'.

Total irrigation was 68.6 mm applied on 3 occasions (R) and 81.0 mm applied on 4 occasions (BB).

**Basal applications:**

Manures (both fields): 1260 kg (0:20:20) plus 900 kg Epsom salts ploughed in in autumn, 360 kg (0:20:20) combine drilled.

**Weedkillers:**

Pastures (R): Paraquat at 0.56 kg ion in 220 l. MCPA, mecoprop and dicamba ('Banlene Plus' at 5.6 l in 220 l).

Windbreak Field (BB): MCPA, mecoprop and dicamba ('Banlene Plus' at 4.2 l) plus mecoprop at 1.4 l in 220 l.

**Seed:** Julia dressed with ethirimol sown at 160 kg.

**Cultivations, etc.:**

Pastures (R): Paraquat applied: 2 Oct, 1972. Basal PK and Epsom salts applied, plots ploughed: 19 Nov. N applied: 17 Mar, 1972. Seed combine drilled: 18 Mar. 'Banlene Plus' applied: 18 May. Irrigated: 13 June, 16 June, 6 July. Combine harvested: 23 Aug. Previous crops: Spring beans 1970, barley 1971.

72/R/B/1 and 72/BB/B/1

Windbreak Field (BB): Basal PK applied: 6 Sept, 1971. Epsom salts applied: 15 Sept. Ploughed: 1 Oct. N applied: 16 Mar, 1972. Seed combine drilled: 17 Mar. Weedkiller applied: 16 May. Irrigated: 2 June, 13 June, 29 June, 30 June. Combine harvested: 11 Aug. Previous crops: Wheat 1970 and 1971.

NOTE: Crop samples were taken throughout the season. The percentage of N, P and K was measured in all samples. The number of tillers or ear-bearing stems was counted, and the leaf areas measured at each sampling. Mildew (*Erysiphe graminis*) assessments were made. 1000 grain weights were determined. Soil moisture content was estimated by neutron probe weekly.

Standard errors per plot. Grain, tonnes/hectare:

Pastures (R): Sub plot: 0.320 or 4.9% (20 d.f.)  
Windbreak Field (BB): Sub plot: 0.414 or 7.5% (20 d.f.)

72/R/B/1 and 72/BB/B/1

TABLES OF MEANS

PASTURES (R)

N: KG/HA

	31	63	94	125	157	188	Mean
GRAIN: TONNES/HECTARE							
I							
O	6.54	6.84	6.95	6.64	6.79	6.49	6.71
C	6.38	6.79	6.60	6.29	5.79	5.86	6.29
Mean	6.46	6.82	6.77	6.47	6.29	6.18	6.50

STANDARD ERRORS OF DIFFERENCES

N	I*
	N
0.185	0.261

STRAW: TONNES/HECTARE

I							
O	5.11	5.73	6.56	6.72	6.50	7.21	6.31
C	5.13	6.51	6.50	7.03	7.09	7.30	6.60
Mean	5.12	6.12	6.53	6.88	6.80	7.25	6.45

Mean D.M. %: Grain: 84.6  
Straw: 86.6

\* Within the same level of I only.

72/R/B/1 and 72/BB/B/1

WINDBREAK FIELD (BB)

N: KG/HA

	31	63	94	125	157	188	Mean
GRAIN: TONNES/HECTARE							
I							
O	4.55	5.13	5.76	5.96	6.19	6.19	5.63
C	4.76	6.10	5.82	5.88	5.24	5.11	5.48
Mean	4.65	5.61	5.79	5.92	5.71	5.65	5.56

STANDARD ERRORS OF DIFFERENCES

N	I* N
0.239	0.338

STRAW: TONNES/HECTARE

I							
O	3.66	5.00	4.80	5.76	5.09	5.73	5.01
C	4.55	5.41	6.83	6.48	6.90	6.03	6.03
Mean	4.10	5.21	5.82	6.12	6.00	5.88	5.52

Mean D.M. %: Grain: 85.1  
Straw: 57.3

\* Within the same level of I only.