

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/S/B/1 Barley Varieties, Ppa, Rates and Times of N

### Rothamsted Research

Rothamsted Research (1973) *72/S/B/1 Barley Varieties, Ppa, Rates and Times of N* ; Yields Of The Field Experiments 1972, pp 320 - 322 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/S/B/1

BARLEY

VARIETIES, PPA, RATES AND TIMES OF N

Object: To study the effects of a range of nitrogen levels, applied to seedbed or in May, on the yield of two barley varieties. The effects of dressing seed with phenylphosphonic acid are also studied - Saxmundham Grove Plot.

Sponsors: F.V. Widdowson, A. Penny.

Design: 2 x 3 x 2 x 2 in 2 randomised blocks of 6 plots split into 2 breadthways and lengthways.

Whole plots dimensions: 2.74 x 12.2. Sub plot area harvested: 0.00052.

Treatments: All combinations of:-

- Whole plots: 1. Varieties: Julia, Midas.  
2. N levels: 50, 100, 150 kg N as calcium nitrate.  
Sub plots: 3. Time of application of N: To seedbed (E), top dressed (L).  
4. Seed dressing (SD): None (O), phenylphosphonic acid (D) at 0.38 kg.

Basal applications: 310 kg (0:20:20) to seedbed. Weedkiller: Dichlorprop plus MCPA ('Mephetol Plus' at 5.6 l). Fungicide: Tridemorph at 0.53 kg.

Seed: Sown at 190 kg.

Cultivations, etc.: Ploughed: Oct, 1971. Seedbed N and basal PK applied, seed drilled: 27 Mar, 1972. N top dressed, weedkiller and fungicide applied: 16 May. Cut by hand: 21 Aug. Previous crop: Barley.

Standard errors per plot. Grain, tonnes/hectare:

Whole plot	0.208 or 4.7% (11 d.f.)
Sub plot Application:	0.410 or 9.3% (12 d.f.)
SD:	0.255 or 5.8% (12 d.f.)
Application x SD:	0.462 or 10.4% (12 d.f.)

72/S/B/1

TABLES OF MEANS

GRAIN: TONNES/HECTARE

VARIETY	N: KG/HA			APPLICATION		SD		Mean
	50	100	150	E	L	O	D	
Julia	3.97	4.90	5.05	4.77	4.51	4.92	4.36	4.64
Midas	4.09	4.28	4.29	4.60	3.84	4.29	4.15	4.22
		N: KG/HA						
			50	4.13	3.94	4.22	3.84	4.03
			100	4.92	4.26	4.76	4.42	4.59
			150	5.01	4.34	4.84	4.50	4.67
				APPLICATION				
				E		4.84	4.53	4.68
				L		4.38	3.97	4.18
Mean						4.61	4.25	4.43

Mean D.M. %: 86.2

STANDARD ERRORS OF DIFFERENCES

VARIETY	N	APPLICATION	SD	VARIETY N	VARIETY APPLICATION	VARIETY SD
0.120	0.147	0.167	0.104	0.208	0.206	0.159
Unless same level of VARIETY					0.237	0.147

	N APPLICATION	N SD	APPLICATION SD
Unless same level of N	0.252	0.195	0.197
APPLICATION SD	0.290	0.181	0.169 0.214



72/S/B/1

STRAW: TONNES/HECTARE

VARIETY	N: KG/HA			APPLICATION		SD		Mean
	50	100	150	E	L	D	D	
Julia	3.69	5.28	6.05	5.26	4.75	5.24	4.77	5.01
Midas	4.81	5.33	6.28	5.81	5.14	5.66	5.28	5.47
			N: KG/HA					
			50	4.33	4.16	4.49	4.01	4.25
			100	5.69	4.92	5.44	5.17	5.31
			150	6.58	5.75	6.42	5.90	6.16
				APPLICATION				
				E		5.74	5.33	5.53
				L		5.16	4.72	4.94
Mean						5.45	5.03	5.24

Mean D.M. %: 86.1