

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1972

[Full Table of Content](#)



72/R/BE/4 Spring Beans Effects of Aphids

Rothamsted Research

Rothamsted Research (1973) *72/R/BE/4 Spring Beans Effects of Aphids* ; Yields Of The Field Experiments 1972, pp 329 - 330 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/BE/4

SPRING BEANS

EFFECTS OF APHIDS

Object: To study the effects of applying liquid or granular insecticides at different times on yield and aphid control on beans - Great Knott III.

Sponsors: R. Bardner, J.H. Stevenson, J.R. Moffatt.

Design: 5 randomised blocks of 6 plots.

Whole plot dimensions: 5.33 x 12.2. Area harvested: 0.00390.

Treatments: Insecticides (I):-

None	(O)
Phorate granules at 1.12 kg:-	
At start of flowering	(GE)
At end of flowering	(GL)
Demeton-s-methyl spray at 0.25 kg in 440 l:-	
At start of flowering	(SE)
At end of flowering	(SL)
At start and again at end of flowering	(SEL)

Basal applications: Ground chalk at 7.5 tonnes. FYM at 30 tonnes. 400 kg (0:14:28) placement drilled. Weedkillers: Paraquat at 0.56 kg ion in 220 l, simazine at 0.84 kg in 220 l.

Seed: Maris Bead, sown at 220 kg.

Cultivations, etc.: Chalked: 3 Sept, 1971. Paraquat applied: 30 Sept. FYM applied: 15 - 22 Nov. Ploughed: 24 Nov. Seed placement drilled: 24 Mar, 1972. Simazine applied: 2 May. Phorate applied: GE plots - 15 June, GL plots - 21 July. Demeton-s-methyl applied: SE and SEL plots - 26 June, SL and SEL plots - 20 July. Combine harvested: 26 Sept. Previous crops: Winter wheat 1970, barley 1971.

NOTE: Aphid counts were made throughout the season. Estimates of plant growth were made. Plant measurements were taken just before harvest.

Corrigendum to 71/R/BE/5: Under 'Treatments' - the amount of demeton-s-methyl sprayed on each occasion was 245 g in 562 l and not as shown.

Standard error per plot.

Grain, tonnes/hectare: 0.356 or 11.6% (20 d.f.)

72/R/BE/4

TABLES OF MEANS

GRAIN: TONNES/HECTARE

I

O	SE	SL	SEL	GE	GL	Mean
3.16	3.11	2.97	3.15	3.19	2.90	3.08

STANDARD ERROR OF DIFFERENCES

I 0.225

Mean D.M. %: 82.1