

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1972

[Full Table of Content](#)



### 72/R/BE/8 Spring Beans N, Carbohydrate and Irrigation

#### Rothamsted Research

Rothamsted Research (1973) *72/R/BE/8 Spring Beans N, Carbohydrate and Irrigation* ; Yields Of The Field Experiments 1972, pp 334 - 335 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/BE/8

SPRING BEANS

N, CARBOHYDRATE AND IRRIGATION

Object: To study the effects of nitrogen fertiliser, sucrose and irrigation, applied at flowering time, on yield, and its components, in field beans - Long Hoos V 5.

Sponsor: J. McEwen.

Design: 3 randomised blocks of 2 plots, split into 4.

Whole plot dimensions: 2.03 x 4.27. Sub plot area harvested: 0.00019.

Treatments: All combinations of:-

Whole plots: 1. Irrigation (I): None (0), irrigated for 3 weeks after first flowers seen (3).

Sub plots: 2. Nitrogen just before flowering: None, 150 kg N as 'Nitro-Chalk'.

3. Carbohydrate (C): None (0), sucrose at 150 kg (half when first flowers seen, half two weeks later) (S).

Irrigation applied by watering can on 16, 17, 24, 27 June, 5 July - 6.3 mm on each occasion. Sucrose was applied as a foliar spray in 1,000 l.

Basal applications: 1570 kg (0:14:28) in autumn. Weedkiller: Simazine at 0.84 kg in 340 l. Insecticide: Demeton-s-methyl at 0.25 kg in 340 l and later at 1 kg in 340 l. Fungicide: Benomyl at 0.75 kg in 1000 l on 2 occasions (applied with sucrose to C plots).

Seed: Minor sown at 224 kg.

Cultivations, etc.: Ploughed: 6 Oct, 1971. Basal PK applied: 30 Sept. Rotary cultivated: 16 Mar, 1972. Seed drilled: 17 Mar. Weedkiller applied: 21 Mar. N applied: 14 June. Benomyl and sucrose applied: 20 June. Insecticide applied: 26 June. Benomyl and sucrose applied: 7 July. Insecticide applied: 19 July. Hand harvested: 28 Sept. Previous crops: Barley 1970, fallow 1971.

NOTE: Counts were made of number of stems and pods. 1000 grain weights and % N in grain were measured.

Standard error per plot.

Grain, tonnes/hectare: Sub plot: 0.157 or 3.4% (12 d.f.).

72/R/BE/8

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	N: KG/HA		C		Mean
	0	150	0	S	
I					
0	4.28	4.64	4.42	4.50	4.46
3	4.61	4.82	4.82	4.60	4.71
		N: KG/HA			
		0	4.47	4.42	4.45
		150	4.78	4.68	4.73
Mean			4.62	4.55	4.59

STANDARD ERRORS OF DIFFERENCES

N	C	I*	I*	N
		N	C	C
0.064	0.064	0.091	0.091	0.091

\* Within the same level of I only

Mean D.M. %: 83.2