

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/R/P/5 Potatoes Seed Sources

### Rothamsted Research

Rothamsted Research (1973) *72/R/P/5 Potatoes Seed Sources* ; Yields Of The Field Experiments 1972, pp 355 - 357 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/P/5

POTATOES

SEED SOURCES

Object: To compare stocks of seed freed from tuber-borne diseases by the use of stem cuttings, with local once-grown and bought-in certified stocks - Delafield.

Sponsors: J.M. Hirst, G.A. Hide, F. Bell.

Design: 2 randomised blocks of 2 plots, split into 24.

Whole plot dimensions: 5.69 x 73.2. Sub plot area harvested: 0.00135.

Treatments: All combinations of:-

Whole plots: 1. Varieties: King Edward (KE), Pentland Crown (PC).

Sub plots: 2. Seed sources:

Virus tested seed from Scotland

(duplicated plots) (1)

Seed once grown at Rothamsted from virus

tested seed from Scotland (duplicated plots) (2)

Certified seed from Scottish or Irish stocks (3-12)

Once grown (English) stocks (13-22)

Basal applications: 1500 kg (13:13:20). Weedkillers: Linuron at 0.84 kg plus paraquat at 0.42 kg ion in 440 l. Fungicide: Mancozeb at 1.3 kg in 440 l on 3 occasions. Insecticide: Demeton-s-methyl at 0.25 kg with the mancozeb on the second occasion.

Cultivations, etc.: Ploughed: 12 Oct, 1971. NPK applied: 20 Apr, 1972.

Rotary cultivated, potatoes machine planted: 21 Apr. Grubbed:

15 June. Earthed up: 16 June. Fungicide applied: 11 July,

24 July, 10 Aug. Insecticide applied: 24 July. Lifted: 2 Oct.

Previous crops: Barley 1970, fallow 1971.

NOTE: Emergence counts were made on May 22. Tubers were graded into 6 sizes and assessments made of Oospora, Rhizoctonia, Helminthosporium and Phoma infection.

Standard error per plot.

Total tubers, tonnes/hectare: Sub plot: 3.54 or 8.6% (50 d.f.)

72/R/P/5

TABLES OF MEANS

TOTAL TUBERS: TONNES/HECTARE

VARIETY

SEED	KE	PC	Mean
1	44.8	42.2	43.5
2	46.7	39.7	43.2
3	42.6	38.4	40.5
4	42.1	40.4	41.2
5	39.0	41.7	40.3
6	42.6	35.5	39.1
7	43.8	34.6	39.2
8	41.7	39.4	40.5
9	43.7	36.5	40.1
10	45.3	41.4	43.4
11	41.5	38.8	40.2
12	42.2	45.2	43.7
13	37.8	40.7	39.3
14	40.6	34.2	37.4
15	41.5	39.3	40.4
16	42.1	38.3	40.2
17	42.1	44.0	43.1
18	43.2	37.3	40.2
19	41.8	42.5	42.1
20	36.8	44.4	40.6
21	46.8	41.3	44.1
22	41.0	41.7	41.3
Mean	42.5	40.0	41.3

STANDARD ERRORS OF DIFFERENCES

SEED

Between 3-22	2.50
1 or 2 v 3-22	2.17
1 v 2	1.77

VARIETY

SEED

Within the same	Between 3-22	3.54
VARIETY	1 or 2 v 3-22	3.07
	1 v 2	2.50



72/R/P/5

PERCENTAGE WARE 4.44 CM (1.75 inch) RIDDLE

SEED	VARIETY		Mean
	KE	PC	
1	61.8	85.4	73.6
2	72.0	85.9	78.9
3	73.9	86.9	80.4
4	58.5	83.9	71.2
5	68.9	86.0	77.5
6	63.0	84.8	73.9
7	72.8	85.7	79.2
8	70.8	86.5	78.6
9	65.9	89.5	77.7
10	74.6	85.8	80.2
11	71.2	84.8	78.0
12	70.8	91.6	81.2
13	65.0	85.5	75.3
14	65.1	91.1	78.1
15	60.5	83.1	71.8
16	64.4	89.1	76.7
17	66.5	85.4	76.0
18	72.0	85.8	78.9
19	74.8	87.0	80.9
20	61.6	88.0	74.8
21	67.1	84.9	76.0
22	66.4	83.7	75.0
Mean	67.6	86.3	76.9