

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1972

[Full Table of Content](#)



72/S/RN/1 Rotation I - Grass, Lucerne

Rothamsted Research

Rothamsted Research (1973) *72/S/RN/1 Rotation I - Grass, Lucerne* ; Yields Of The Field Experiments 1972, pp 41 - 44 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/S/RN/1

ROTATION I

Object: To compare nutrient cycles, uptakes of nutrients and responses to fresh P and K of lucerne and grass leys. To obtain an estimate of the rate of release of nutrients, particularly K, from Saxmundham soil. The effects of lucerne and grass leys will be compared on subsequent arable crops - Saxmundham.

Sponsors: A.E. Johnston, R.C. Flint.

For previous years see 'Details' 1967, 68/A/9(t), 69/S/RN/1(t), 70/S/RN/1(t) and 71/S/RN/1.

Whole plot dimensions (new plots): 5.49 x 17.1.

Area harvested: Grass: 1st cut: New treatments: 0.00003
Old treatments: 0.00050
2nd cut: New treatments: 0.00139
Old treatments: 0.00050
Lucerne: New treatments: 0.00106
Old treatments: 0.00139

Seed:- Grass: Timothy S352 and Meadow Fescue S215
Lucerne: Europe.

Cultivations, etc.:

Grass: P and K applied: 15 Feb, 1972. Bonemeal applied: 22 Feb.
N applied: 27 Mar. Cut: Blocks 3 and 4 - 23 May, 16 Aug,
blocks 1 and 2 - 1 June, 16 Aug. N applied after the first
cut.

Lucerne: P and K applied: 15 Feb, 1972. Bonemeal applied:
22 Feb. Cut: 7 June, 10 Aug.

72/S/RN/1

TABLES OF MEANS

OLD TREATMENTS

GRASS, DRY MATTER: TONNES/HECTARE

Treatment 1899-1972	1st cut	2nd cut	Total of 2 cuts
D*	2.38	0.53	2.92
B	0.61	0.00	0.61
N	2.68	1.19	3.86
P	0.69	0.00	0.69
K	0.51	0.00	0.51
-	0.58	0.00	0.58
PK	0.72	0.00	0.72
NK	2.41	1.44	3.85
NP	2.52	1.49	4.01
NPK	2.93	1.93	4.87
Mean	1.60	0.66	2.26

Mean D.M. %: 1st cut: 27.7
 2nd cut: 24.1
 Total of 2 cuts: 25.9

* Last applied autumn 1969

72/S/RN/1

NEW TREATMENTS

GRASS, DRY MATTER: TONNES/HECTARE

Treatment 1899-1965	Treatment* from 1966	1st cut	2nd cut	Total of 2 cuts
D	DN1	6.81	3.88	10.69
B	B	5.41	3.71	9.12
N	N2P2	5.90	3.58	9.48
P	N1P1	5.60	3.79	9.39
K	N1P2K	5.82	3.99	9.81
-	N1P2	5.52	3.65	9.17
PK	N1P1K	5.89	4.08	9.97
NK	N2P2K	5.95	4.09	10.04
NP	N2P1	5.63	3.51	9.14
NPK	N2P1K	6.10	3.88	9.98
Mean		5.86	3.82	9.68

Mean D.M. %: 1st cut: 22.2
 2nd cut: 43.1
 Total of 2 cuts: 32.6

* N not tested since 1970 - all treatments receive basal N at 100 kg for each cut. FYM last applied autumn 1969.

72/S/RN/1

NEW TREATMENTS

LUCERNE, DRY MATTER: TONNES/HECTARE

Treatment 1899-1965	Treatment* from 1966	1st cut	2nd cut	Total of 2 cuts
D	DN1	5.37	5.33	10.70
B	B	3.96	4.54	8.50
N	N2P2	4.29	4.63	8.92
P	N1P1	3.87	4.63	8.50
K	N1P2K	4.73	5.33	10.06
-	N1P2	4.08	4.70	8.78
PK	N1P1K	4.61	5.17	9.77
NK	N2P2K	4.64	5.08	9.72
NP	N2P1	3.82	4.40	8.22
NPK	N2P1K	4.36	5.07	9.43
Mean		4.37	4.89	9.26

Mean D.M. %: 1st cut: 32.2
 2nd cut: 27.0
 Total of 2 cuts: 29.6

* N not applied since 1970. FYM last applied autumn 1969.