

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/R/RN/8 Cultivation/WEEDKILLER - Beans, Wheat, Potatoes, Barley

### Rothamsted Research

Rothamsted Research (1973) 72/R/RN/8 Cultivation/WEEDKILLER - Beans, Wheat, Potatoes, Barley ;  
Yields Of The Field Experiments 1972, pp 92 - 98 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/RN/8

CULTIVATION/WEEDKILLER

Object: To determine the long-term effects of weedkillers and different methods of primary cultivation on a rotation of crops - Great Harpenden I.

Sponsors: J.R. Moffatt, G.V. Dyke, J.A. Currie.

The 12th year, beans, wheat, potatoes, barley.

For previous years see 'Details' 1967, 68/B/6(t), 69/R/RN/8(t), 70/R/RN/8 and 71/R/RN/8(t).

Whole plot dimensions: 12.8 x 15.2. Area harvested: Beans: 0.00488, winter wheat and barley: 0.00434, potatoes: 0.00217.

Seed: Beans: Maris Bead sown at 220 kg.

Wheat: Cappelle sown at 200 kg.

Potatoes: Pentland Crown, Rothamsted once grown.

Barley: Julia, seed dressed ethirimol, sown at 160 kg.

Cultivations, etc.:-

Spring beans: Barley straw burnt on B plots: 25 Aug, 1971. Paraquat applied at 0.56 kg ion in 220 l to G sub plots: 10 Sept. B plots deep-tine cultivated once, T plots twice: 4 Nov. R plots rotary cultivated, P and C plots ploughed: 8 Nov. P, R, T, B and C plots power-harrowed, A plots rotary cultivated: 20 Mar, 1972. All plots placement drilled: 21 Mar. Simazine applied at 1.1 kg in 220 to SX, A, B and C plots: 2 May. M plots tractor hoed: 4 May. M plots mechanically weeded: 9 May. Dinoseb acetate ('Ivosit' at 6.7 kg) applied in 440 l to SY plots: 19 May. M plots tractor hoed: 25 May, 7 June. Phorate applied at 1.1 kg: 16 June. Ends of all plots sprayed with demeton-s-methyl at 0.25 kg in 440 l: 5 Aug. Combine harvested: 26 Sept.

Wheat: Bean straw raked off B plots (insufficient to burn): 6 Sept, 1971. Paraquat applied at 0.56 kg ion in 220 l to G sub plots: 10 Sept. B plots deep-tine cultivated once, T plots twice: 8 Oct. P, A and C plots ploughed: 11 Oct. P, T, A, B and C plots spring-tine cultivated, R plots rotary cultivated, seed combine drilled: 15 Oct. All plots harrowed: 22 Mar, 1972. All plots rolled: 25 Mar. N applied: 19 Apr. H sub plots sprayed 0.63 kg ioxynil and 1.9 kg mecoprop in 220 l: 28 Apr. Combine harvested: 31 Aug.

72/R/RN/8

Potatoes: Wheat straw burnt on B plots: 7 Sept, 1971. Paraquat applied to G sub plots: 10 Sept. T plots deep-tine cultivated twice: 4 Nov. P and C plots ploughed, R plots rotary cultivated: 8 Nov. Basal NPK applied, P and T plots spring-tine cultivated: 18 Apr, 1972. P and T plots spring-tine cultivated: 19 Apr. R, A, B and C plots rotary cultivated, potatoes machine planted: 19 - 20 Apr. All ridges rolled: 2 May. M plots chain harrowed and grubbed: 9 May. S plots sprayed linuron at 0.84 kg and paraquat at 0.42 kg ion in 440 l: 17 May. M and Y plots grubbed: 14 June. M and Y plots rotary ridged: 15 June. Sprayed three times with mancozeb at 1.34 kg in 440 l: 11 July, 21 July, 10 Aug. Demeton-s-methyl at 0.25 kg was included with the second spray. Sprayed with undiluted BOV at 200 l: 20 Sept. Lifted: 7 - 9 Nov.

Barley: Spring-tine cultivated: 12 Oct, 1971. Sodium trichloroacetate applied at 20 kg in 220 l: 21 Oct. Spring-tine cultivated twice: 28 Oct, 17 Nov. Sodium trichloroacetate applied at 20 kg in 220 l: 2 Dec. Spring-tine cultivated: 10 Dec. T and C plots deep-tine cultivated twice, R plots rotary cultivated, P plots ploughed: 13 Dec. A plots rotary cultivated, P, R, T, B and C plots power-harrowed: 20 Mar, 1972. Seed combine drilled: 21 Mar. Rolled: 22 Mar. H sub-plots sprayed 0.53 kg ioxyail and 1.6 kg mecoprop in 220 l: 17 May. Combine harvested: 18 Aug.

- NOTES: (1) Docks were hand pulled in potatoes and thistles in all crops. Weed counts were taken.
- (2) Two rows of potatoes per sub-plot were lifted for yield by elevator digger (not 4 as in previous years), then the 2 adjacent rows were lifted by harvester. Yields are presented for the elevator digger only.

Standard errors per plot.

Spring beans: Grain, tonnes/hectare:	Whole plot: 0.298 or 9.3% (8 d.f.)
	Sub plot: 0.344 or 10.8% (9 d.f.)
Spring wheat: Grain, tonnes/hectare:	Whole plot: 0.362 or 5.4% (8 d.f.)
	Sub plot: 0.291 or 4.4% (8 d.f.)
Potatoes, Total tubers, tonnes/hectare:	Whole plot: 3.48 or 9.4% (8 d.f.)
	Sub plot: 2.74 or 7.4% (9 d.f.)
Barley: Grain, tonnes/hectare:	Whole plot: 0.164 or 2.4% (8 d.f.)
	Sub plot: 0.275 or 4.0% (9 d.f.)

72/R/RN/8

TABLES OF MEANS

SPRING BEANS

GRAIN: TONNES/HECTARE

CULTIVATION (C)

	P	R	T	Mean
Mean	3.47	2.99	3.12	3.19
WEEDKILLER (W)				
M	3.64	2.97	3.37	3.33
SX	3.10	2.95	2.64	2.90
SY	3.68	3.05	3.34	3.35
PARAQUAT SPRAY (P)				
O	3.52	2.89	3.01	3.14
G	3.42	3.09	3.22	3.24

STANDARD ERRORS OF DIFFERENCES

	C	W	P	C W	C P
Unless same level of	0.172	0.172	0.115	0.298	0.222
C					0.199

A	AG	BG	C	CG
2.95	2.93	2.94	3.17	3.41

Grand mean: 3.16  
Mean D.M. %: 83.2

72/R/RN/8

WINTER WHEAT

GRAIN: TONNES/HECTARE

CULTIVATION (C)

	P	R	T	Mean
Mean	7.07	6.56	6.40	6.68
WEEDKILLER 1971 (W)				
M	6.98	6.50	6.58	6.68
SX	7.07	7.06	6.53	6.88
SY	7.17	6.11	6.09	6.46
PARAQUAT SPRAY 1972 (P)				
O	6.94	6.38	6.18	6.50
G	7.21	6.73	6.62	6.85
HORMONE SPRAY 1972 (H)				
O	6.49	6.65	6.41	6.52
H	6.88	7.12	6.51	6.83

STANDARD ERRORS OF DIFFERENCES

	C	W	P	H	C W	C P	C H
Unless same level of C	0.209	0.209	0.097	0.097	0.362	0.241	0.241
						0.168	0.168

A	AG	AH	AGH	BGH	CH	CGH
7.35	6.93	6.56	7.57	7.12	7.52	6.67

Grand mean: 6.78  
Mean D.M. %: 84.5

72/R/RN/8

POTATOES

TOTAL TUBERS: TONNES/HECTARE

CULTIVATION (C)

	P	R	T	Mean
Mean	39.5	34.7	37.1	37.1
WEEDKILLER (W)				
M	42.3	34.0	34.5	37.0
S	36.4	35.7	34.4	35.5
SY	39.8	34.6	42.3	38.9
PARAQUAT SPRAY (P)				
O	37.3	35.3	35.7	36.1
G	41.7	34.2	38.5	38.1

STANDARD ERRORS OF DIFFERENCES

	C	W	P	C W	C P
Unless same level of C	2.01	2.01	0.91	3.48	2.30
					1.58

A	AG	BG	C	CG
32.1	32.4	33.7	43.6	40.7

Grand mean: 36.8

72/R/RN/8

POTATOES

PERCENTAGE WARE: 3.81 CM (1.5 INCH) RIDDLE

CULTIVATION (C)

	P	R	T	Mean
Mean	95.0	96.2	95.8	95.7
WEEDKILLER (W)				
M	96.8	97.3	95.5	96.6
S	93.2	95.2	95.3	94.6
SY	95.1	96.0	96.5	95.9
PARAQUAT SPRAY (P)				
O	94.6	96.1	95.5	95.4
G	95.5	96.3	96.1	96.0

A	AG	BG	C	CG
97.3	96.2	94.0	95.6	95.1

Grand mean: 95.6

72/R/RN/8

BARLEY

GRAIN: TONNES/HECTARE

CULTIVATION (C)

	P	R	T	Mean
Mean	6.88	6.82	6.80	6.83
WEEDKILLER 1971 (W)				
M	6.87	6.95	6.85	6.89
S	6.81	6.70	6.56	6.69
SY	6.95	6.82	7.00	6.92
HORMONE SPRAY 1972 (H)				
O	6.89	6.81	6.67	6.79
H	6.87	6.84	6.93	6.88

STANDARD ERRORS OF DIFFERENCES

	C	W	H	C W	C H
Unless same level of	0.095	0.095	0.092	0.164	0.147
C					0.158
A	6.31				
AH		6.64			
BH			6.78		
CH				6.93	

Grand mean: 6.81  
 Mean D.M. %: 82.7