

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1973

[Full Table of Content](#)



73/R/CS/42 - Effects of Invertebrates on Yield - Old Grass

Rothamsted Research

Rothamsted Research (1974) *73/R/CS/42 - Effects of Invertebrates on Yield - Old Grass* ; Yields Of The Field Experiments 1973, pp 206 - 208 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/R/CS/42

EFFECT OF INVERTEBRATES ON YIELD

Object: To study the effects of a range of invertebrate-killing chemicals on the yield of old grass - Road Piece.

Sponsor: I.F. Henderson.

The fifth year, old grass.

For previous years see 69/R/CS/42(t), 70/R/CS/42(t), 71/R/CS/42(t) and 72/R/CS/42(t).

Design: 4 randomised blocks of 12 plots.

Whole plot dimensions: 2.74 x 6.40. Area harvested: 0.00065.

Treatments: Chemicals:	CHEMICAL
None	0
To control foliar invertebrates:	
Chlorbenside spray until 1972, none since	(FCB)
Menazon spray, repeated frequently	FMZ
Chlorbenside (until 1972), menazon, dimethoate sprays and parathion/bran bait, repeated frequently	F3
To control soil invertebrates:	
Formalin, last applied in 1970	(SFO)
Parathion/bran bait, repeated frequently	SL
Nematicide C14421, last applied in 1970	(SN)
Metalddehyde, repeated frequently	SMT
Aldrin in 1969 and 1972 only, nematicide C14421 until 1970, metalddehyde repeated frequently	S3
As S3 + chlordane in 1969 and 1972	S4
To control foliar and soil invertebrates:	
As F3+S4 since 1969	F3S4
As F3+S4 since 1972 (previously untreated)	F3S4*

73/R/CS/42

Basal applications: Manures: (0:14:28) at 500 kg in winter, (25:0:16) at 440 kg in spring, (25:0:16) at 220 kg after each cut except the last.

Cultivations, etc.: - PK applied: 17 Nov, 1972. NK applied: 21 Feb, 1973. Cut three times: 22 May, 4 July, 12 Sept. NK applied after the first two cuts.

Chemical treatments applied:

Dimethoate: 11 May, 22 June, 23 July, 16 Aug.

Menazon: 10 May, 21 June, 20 July, 16 Aug.

Metaldehyde: 10 May, 21 June, 24 July, 15 Aug.

Parathion: 11 May, 21 June, 24 July, 25 Aug.

NOTE: Samples were taken for botanical analysis in May. Samples for fauna were taken after the last cut.

Standard errors per plot. Dry matter, tonnes/hectare:

1st cut: 0.505 or 11.3% (33 d.f.)

2nd cut: 0.305 or 15.6% (33 d.f.)

3rd cut: 0.234 or 11.6% (33 d.f.)

Total of 3 cuts: 0.586 or 6.9% (33 d.f.)

73/R/CS/42

TABLES OF MEANS

DRY MATTER: TONNES/HECTARE

	1ST CUT	2ND CUT	3RD CUT	TOTAL OF 3 CUTS
CHEMICAL				
0	3.91	1.96	2.26	8.13
(FCB)	3.75	2.14	2.12	8.01
FMZ	4.57	1.99	2.00	8.57
F3	4.95	1.80	1.66	8.40
(SFD)	4.20	1.73	2.41	8.34
SL	4.62	1.86	2.27	8.75
(SN)	4.16	1.97	2.24	8.37
SMT	4.08	2.14	1.87	8.10
S3	4.68	1.97	2.02	8.67
S4	4.58	2.07	1.99	8.63
F3S4	5.38	1.65	1.35	8.39
F3S4*	4.95	2.13	2.05	9.12
Mean	4.49	1.95	2.02	8.46

STANDARD ERRORS OF DIFFERENCES

CHEMICAL	0.357	0.216	0.166	0.415
Mean D.M. %	1st cut:	16.8		
	2nd cut:	23.4		
	3rd cut:	36.2		
	Total of 3 cuts:	25.5		

* Since 1972