

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1973

[Full Table of Content](#)



---

### 73/W/CS/66 - Dazomet & Nitrogen - Maize

#### Rothamsted Research

Rothamsted Research (1974) 73/W/CS/66 - *Dazomet & Nitrogen - Maize* ; Yields Of The Field Experiments 1973, pp 219 - 220 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/W/CS/66

DAZOMET AND NITROGEN

Object: To study the cumulative effects of dazomet and nitrogen on pathogens and yield of maize grown continuously - Woburn Butt Furlong.

Sponsors: A.J. Barnard, D. Hornby.

The third year, maize.

For previous years see 71/W/CS/66(t) and 72/W/CS/66(t).

Design: 4 blocks of 2 plots split into 4.

Whole plot dimensions: 2.13 x 16.5. Area harvested: 0.00053.

Treatments: All combinations of:-

Whole plots: 1. Dazomet (kg per annum) cumulative 1971-73: DAZOMET

0	0
450	450

Sub plots: 2. Nitrogen fertiliser (kg N per annum) cumulative 1971-73:

	N
50 to seedbed	50
100 to seedbed	100
150 to seedbed	150
100 to seedbed, 50, five weeks after germination	100+50

Basal applications: Manures: (0:14:28) at 870 kg. Weedkiller: Atrazine at 1.1 kg in 280 l.

Seed: Pioneer 131, sown at 30 kg.

Cultivations, etc.:- Deep-tine cultivated: 11 Aug, 1972. Dazomet applied, all plots rotary cultivated twice: 12 Oct. Rolled: 13 Oct. Ploughed: 3 Jan, 1973. PK applied, weedkiller applied and harrowed in: 27 Apr. N applied, seed sown: 2 May. Late N applied: 2 July. Harvested by hand: 1 Nov.

73/W/CS/66

- NOTES: (1) Soil samples were taken in spring before sowing and again after harvest for counts of ectoparasitic nematodes.  
 (2) Leaf samples were taken in September for analysis of % P and K.  
 (3) Plant samples were taken in September for incidence of stem and leaf pathogens.  
 (4) % N in grain was determined.  
 (5) Two plots, DAZOMET 450, N 150 and N 50, were waterlogged and one DAZOMET 0, N 100+50 was damaged by birds. Estimated values were used in the analysis.

Standard error per sub plot.

Grain, tonnes/hectare: 0.835 or 14.5% (15 d.f.)

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	N				Mean
	50	100	150	100+50	
DAZOMET					
0	5.33	4.55	5.66	6.08	5.41
450	5.88	6.46	6.05	5.92	6.08
Mean	5.60	5.51	5.85	6.00	5.74

STANDARD ERRORS OF DIFFERENCES

N	DAZOMET*
N	
0.417	0.590

\* Within the same level of DAZOMET only

Mean D.M. % 64.7