


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1973

[Full Table of Content](#)



73/R&W/CS/88 - Fertiliser & Fym - Wheat

Rothamsted Research

Rothamsted Research (1974) 73/R&W/CS/88 - *Fertiliser & Fym - Wheat* ; Yields Of The Field Experiments 1973, pp 235 - 240 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/R/CS/88 and 73/W/CS/88

FERTILISER AND FYM

Object: To study the residual effects of a range of rates of NPK fertiliser and FYM applied to potatoes on the yield of winter wheat - Rothamsted, Great Knott I (R) and Woburn, Warren Field I (W).

Sponsor: F.V. Widdowson.

The second year, winter wheat.

For previous year see 72/RW/CS/88(t).

Design: 3 randomised blocks of 18 plots.

Whole plot dimensions: 4.27 x 16.2. Area harvested: 0.00217.

Treatments: All combinations of:-

Whole plots: 1. Farmyard manure to potatoes in 1972 (tonnes): FYM

(R)	(W)	(R)	(W)
None	None	0	0
53	73	53	73

2. N and PK fertilisers to potatoes in 1972 to give rates of nitrogen* (kg N): NRESID

188	188
377	377
565	565

3. Times of applying PK fertilisers to potatoes: TIME

All in autumn 1971	Autumn
All in spring 1972	Spring
Half in autumn, half in spring	Aut/Spng

Sub plots: 4. Nitrogen fertiliser to wheat in 1973 (kg N): N

None	0
63	63

* the ratio of N:P2O5:K2O was 1:1.5:1.5 for all NRESID treatments.

Basal applications:

Great Knott I (R): Weedkiller: Dicamba, mecoprop and MCPA ('Tetralex Plus' at 7 l in 220 l).

73/R/CS/88 and 73/W/CS/88

2

Warren Field I (W): Weedkiller: Ioxynil at 0.63 kg plus mecoprop at 1.9 kg
in 280 l.

Seed: Great Knott I (R): Cappelle, sown at 200 kg.
Warren Field I (W): Cappelle, sown at 190 kg.

Cultivations, etc.: -

Great Knott I (R): Deep-tine cultivated twice: 23 Oct, 1972. Power
harrowed: 25 Oct. Seed sown: 26 Oct. N applied: 13 Apr, 1973.
Weedkiller applied: 26 Apr. Combine harvested: 22 Aug.
Warren Field I (W): Deep-tine cultivated: 19 Oct, 1972. Seed sown
24 Oct. N applied: 19 Apr, 1973. Weedkiller applied: 26 Apr.
Combine harvested: 22 Aug.

Standard errors per plot. Grain, tonnes/hectare.

Great Knott I (R): Whole plot: 0.429 or 7.0% (34 d.f.)
Sub plot: 0.688 or 11.2% (36 d.f.)
Warren Field I (W): Whole plot: 0.385 or 8.1% (34 d.f.)
Sub plot: 0.471 or 9.9% (36 d.f.)

73/R/CS/88 and 73/W/CS/88

TABLES OF MEANS

GREAT KNOTT I (R)

GRAIN: TONNES/HECTARE

	NRESID			TIME			N		Mean
	188	377	565	Autumn	Spring	Aut/Sprng	0	63	
FYM									
0	7.26	5.95	5.62	6.16	6.12	6.55	6.44	6.11	6.27
53	6.69	5.59	5.66	5.79	6.07	6.07	6.17	5.78	5.98
	NRESID								
	188	6.89	6.82	7.21	7.21	6.74	6.97		
	377	5.53	5.84	5.93	5.96	5.58	5.77		
	565	5.50	5.62	5.79	5.75	5.52	5.64		
	TIME								
		Autumn		6.13	5.82	5.97			
		Spring		6.35	5.84	6.09			
		Aut/Sprng		6.44	6.18	6.31			
Mean							6.31	5.95	6.13

STANDARD ERRORS OF DIFFERENCES

FYM	NRESID	TIME	N	FYM	NRESID	TIME	FYM	NRESID	TIME
							N	N	N
0.117	0.143	0.143	0.132	0.202	0.202	0.248	0.177	0.216	0.216
Except when comparing means with same level of									
FYM							0.187		
NRESID								0.229	
TIME									0.229

Mean D.M. % 83.6

73/R/CS/88 and 73/W/CS/88

GREAT KNUTT I (R)

STRAW: TONNES/HECTARE

FYM	NRESID			TIME			N		Mean
	188	377	565	Autumn	Spring	Aut/Spng	0	63	
0	7.11	9.28	9.22	8.56	8.36	8.69	8.34	8.73	8.54
53	8.73	9.38	9.39	9.30	9.01	9.18	9.07	9.26	9.16
NRESID				188	8.18	7.45	8.11	7.35	8.48
				377	9.21	9.28	9.51	9.28	9.38
				565	9.40	9.32	9.19	9.49	9.12
				TIME					
				Autumn			8.74	9.12	8.93
				Spring			8.56	8.81	8.69
				Aut/Spng			8.82	9.05	8.93
Mean							8.71	8.99	8.85

Mean D.M. % 84.0

73/R/CS/88 and 73/W/CS/88

WARREN FIELD I (W)

GRAIN: TONNES/HECTARE

	NRESID			TIME			N		
	188	377	565	Autumn	Spring	Aut/Spng	0	63	Mean
FYM									
0	5.35	5.25	3.71	4.75	5.00	4.55	4.84	4.70	4.77
73	5.68	4.84	3.87	4.56	4.82	5.01	5.02	4.57	4.80
	NRESID								
	188	5.66	5.43	5.46	4.94	6.09	5.52		
	377	4.60	5.40	5.13	5.58	4.50	5.04		
	565	3.71	3.92	3.74	4.26	3.32	3.79		
	TIME								
		Autumn		4.83	4.48	4.66			
		Spring		5.03	4.80	4.91			
		Aut/Spng		4.92	4.64	4.78			
Mean							4.93	4.64	4.78

STANDARD ERRORS OF DIFFERENCES

FYM	NRESID	TIME	N	FYM	NRESID	TIME	FYM	NRESID	TIME
								N	N
0.105	0.128	0.128	0.091	0.182	0.182	0.222	0.139	0.170	0.170
Except when comparing means with same level of									
FYM							0.128		
NRESID								0.157	
TIME									0.157

Mean D.M. % 83.2

73/R/CS/86 and 73/W/CS/86

WARREN FIELD I (W)

STRAW: TONNES/HECTARE

FYM	NRESID			TIME			0	63	Mean
	188	377	565	Autumn	Spring	Aut/Spring			
0	5.38	6.71	6.69	6.35	6.62	6.31	6.12	6.73	6.43
73	6.09	6.93	6.48	6.54	6.43	6.54	6.57	6.43	6.50
NRESID									
	188	6.38	5.76	5.81	5.04	6.93	5.98		
	377	6.54	6.74	7.18	7.05	6.59	6.82		
	565	6.41	7.08	6.28	6.95	6.22	6.59		
TIME									
		Autumn		6.25	6.64	6.44			
		Spring		6.38	6.67	6.52			
		Aut/Spring		6.41	6.44	6.42			
Mean					6.35	6.58	6.46		

Mean D.M. % 88.4