

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1973

[Full Table of Content](#)



### 73/R&W/CS/110 - Fertiliser I Fym - Potatoes

#### Rothamsted Research

Rothamsted Research (1974) 73/R&W/CS/110 - *Fertiliser I Fym - Potatoes* ; Yields Of The Field Experiments 1973, pp 271 - 276 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/R/CS/110 and 73/W/CS/110

FERTILISER AND FYM

Object: To study the effects of a range of rates of NPK fertiliser, P and K being applied in autumn or spring, and FYM on the yields of potatoes - Rothamsted, Stackyard (R) and Woburn, Great Hill II (W).

Sponsor: F.V. Widdowson.

The first year, potatoes.

Design: 3 randomised blocks of 18 plots.

Whole plot dimensions: 4.27 x 16.2. Area harvested: 0.00230.

Treatments: All combinations of:-

1. Farmyard manure (tonnes) to supply 377 kg N:	FYM
None	0 0
80 Stackyard (R), 63 Great Hill II (W)	80 63
2. N, in spring and PK fertilisers to give rates of nitrogen* (kg N):	N
188	188
377	377
565	565
3. Times of applying PK fertilisers	TIME
All in autumn	Autumn
All in spring	Spring
Half in autumn, half in spring	Aut/Spng

\* The ratio of N:P2O5:K2O was 1:1.5:1.5 for all N treatments.

Basal applications:

Stackyard (R): Weedkiller: Linuron at 1.2 kg plus paraquat at 0.42 kg ion in 450 l. Fungicide with insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 370 l. Fungicide: Mancozeb at 1.3 kg in 370 l.

Great Hill II (W): Manures: Magnesian limestone at 7.5 tonnes. Weed-killers: Linuron at 1.2 kg plus paraquat at 0.56 kg ion in 280 l. Fungicide with insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 390 l. Fungicide: Mancozeb at 1.3 kg in 390 l on the first occasion and in 370 l on the second occasion.

73/R/CS/110 and 73/W/CS/110

Seed: Stackyard (R): King Edward,  
Great Hill II (W): Maris Piper.

Cultivations, etc.:-

Stackyard (R): Autumn PK applied: 31 Oct, 1972. FYM applied, ploughed: 3 Nov. Spring PK applied: 28 Mar, 1973. N applied, rotary cultivated, potatoes planted: 11 Apr. Weedkiller applied: 15 May. Grubbed: 7 June. Rotary ridged: 11 June. Fungicide with insecticide applied: 2 July. Fungicide applied: 18 July, 9 Aug. Haulm mechanically destroyed: 17 Sept. Sprayed with undiluted BCV at 220 l: 18 Sept. Lifted: 2 Oct. Previous crops: Wheat 1971, fallow 1972.

Great Hill II (W): Magnesian limestone applied: 19 Sept, 1972. Autumn PK applied: 16 Nov. FYM applied, ploughed: 29 Nov. Spring PK applied: 29 Mar, 1973. N applied: 3 Apr. Rotary cultivated, potatoes planted: 6 Apr. Weedkiller applied: 9 May. Grubbed: 31 May. Rotary ridged: 5 June. Fungicide with insecticide applied: 6 July. Fungicide applied: 26 July, 13 Aug. Haulm mechanically destroyed: 12 Sept. Lifted: 14 Sept. Previous crops: Potatoes 1971, winter wheat 1972.

- NOTES: (1) Leaf samples were taken in July for percentage of Mg and K.  
(2) Tuber samples were taken at grading for percentage of N, P and K.  
(3) Tuber samples were taken for cooking tests by N.I.A.B.

Standard errors per plot. Total tubers, tonnes/hectare:

Stackyard (R): 2.75 or 5.4% (34 d.f.)

Great Hill II (W): 2.82 or 7.2% (34 d.f.)

73/R/CS/110 and 73/W/CS/110

TABLES OF MEANS

STACKYARD (R)

TOTAL TUBERS: TONNES/HECTARE

	188	N		TIME			Mean
		377	565	Autumn	Spring	Aut/Spng	
FYM							
0	50.2	49.6	49.2	49.7	48.9	50.4	49.6
80	55.3	52.8	49.6	53.1	51.8	52.8	52.6
		N					
		188		51.1	54.2	53.0	52.7
		377		52.8	48.0	52.8	51.2
		565		50.3	48.9	48.9	49.4
Mean				51.4	50.3	51.6	51.1

TIME	188			377			565		
	Autumn	Spring	Aut/Spng	Autumn	Spring	Aut/Spng	Autumn	Spring	Aut/Spng
FYM									
0	48.0	51.9	50.8	51.4	45.5	51.8	49.8	49.2	48.5
80	54.2	56.4	55.2	54.1	50.5	53.8	50.8	48.5	49.4

STANDARD ERRORS OF DIFFERENCES

FYM	N	TIME	FYM N	FYM TIME	N TIME	FYM N TIME
0.75	0.92	0.92	1.30	1.30	1.59	2.24



73/R/CS/110 and 73/W/CS/110

STACKYARD (R)

PERCENTAGE WARE: 4.44 CM (1.75 INCH) RIDDLER

	188	N		TIME			Mean
		377	565	Autumn	Spring	Aut/Spng	
FYM							
0	78.9	83.4	81.7	84.2	79.9	79.8	81.3
80	83.8	81.8	83.2	84.4	82.3	82.1	82.9
		N					
		188		83.8	80.4	79.8	81.3
		377		85.0	80.3	82.5	82.6
		565		84.1	82.6	80.7	82.5
Mean				84.3	81.1	81.0	82.1

N TIME	188			377			565		
	Autumn	Spring	Aut/Spng	Autumn	Spring	Aut/Spng	Autumn	Spring	Aut/Spng
FYM									
0	82.2	77.8	76.7	85.2	80.3	84.7	85.4	81.7	78.0
80	85.4	83.0	82.9	84.8	80.2	80.2	82.9	83.5	83.3

73/R/CS/110 and 73/W/CS/110

GREAT HILL II (W)

TOTAL TUBERS: TONNES/HECTARE

	188	N		565	TIME			Mean
		377			Autumn	Spring	Aut/Spng	
FYM								
0	33.2	40.9	41.4	36.7	39.6	39.3	38.5	
63	35.1	43.3	42.1	37.6	41.7	41.1	40.2	
		N						
		188		31.8	35.9	34.7	34.1	
		377		38.9	43.2	44.3	42.1	
		565		40.8	42.8	41.5	41.7	
Mean				37.2	40.6	40.2	39.3	

N TIME	188			377			565		
	Autumn	Spring	Aut/Spng	Autumn	Spring	Aut/Spng	Autumn	Spring	Aut/Spng
FYM									
0	30.4	34.2	35.0	38.2	41.9	42.8	41.6	42.5	40.0
63	33.1	37.6	34.4	39.6	44.5	45.9	40.1	43.0	43.0

STANDARD ERRORS OF DIFFERENCES

FYM	N	TIME	FYM N	FYM TIME	N TIME	FYM N TIME
0.77	0.94	0.94	1.33	1.33	1.63	2.30

73/R/CS/110 and 73/W/CS/110

GREAT HILL II (W)

PERCENTAGE WARE: 4.44 CM (1.75 INCH) RIDDLER

	188	N		TIME			Mean
		377	565	Autumn	Spring	Aut/Spng	
FYM							
0	53.2	66.6	68.7	60.2	65.0	63.2	62.8
63	58.9	67.1	67.3	62.4	65.4	65.5	64.4
		N					
		188		53.2	58.5	56.5	56.1
		377		64.5	68.6	67.3	66.8
		565		66.2	68.4	69.3	68.0
Mean				61.3	65.2	64.3	63.6

N TIME	188			377			565		
	Autumn	Spring	Aut/Spng	Autumn	Spring	Aut/Spng	Autumn	Spring	Aut/Spng
FYM									
0	49.9	54.3	55.3	63.0	69.3	67.4	67.8	71.5	66.8
63	56.5	62.7	57.7	66.1	68.0	67.1	64.7	65.4	71.8