

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1975

[Full Table of Content](#)



## 75/ W/RN/13 - Intensive Cereals - Ley, Potatoes, Wheat, Barley

### Rothamsted Research

Rothamsted Research (1976) 75/ W/RN/13 - *Intensive Cereals - Ley, Potatoes, Wheat, Barley* ; Yields Of The Field Experiments 1975, pp 113 - 117 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/W/RN/13

INTENSIVE CEREALS

Object: To study the effects of intensive cereal cropping on yield, incidence of soil-borne diseases and organic matter in the soil - Woburn Stackyard I.

Sponsor: D.B. Slope.

The tenth year, ley, potatoes, winter wheat, barley.

For previous years see 66/B/9(t), 67/B/9, 68/B/7(t), 69/W/RN/13(t), 70/W/RN/13(t), 71/W/RN/13(t) and 72-74/W/RN/13.

Design: For each experiment: 2 randomised blocks of 6 plots, split into 4.

Whole plot dimensions: 8.53 x 20.4.

Treatments:-

One experiment on winter wheat on part of the site of the classical wheat experiment 1877-1954

One experiment on barley on part of the site of the classical barley experiment 1877-1954

Factors tested on both experiments are the same but crop and nitrogen rates differ. All combinations of:-

Whole plots: 1. Previous crops:

PREVCROP

1968 1969 1970 1971 1972 1973 1974

L	P	C	C	C	L	P	C/C/L/P
P	C	C	C	L	P	C	C/L/P/C
C	C	C	L	P	C	C	L/P/C/C
C	C	L	P	C	C	C	P/C/C/C
C	L	P	C	C	C	L	C/C/C/L
C	C	C	C	C	C	C	C/C/C/C

Ley = 1 year ley P = Potatoes C = Cereal: wheat or barley.

Sub plots: 2. Nitrogen fertiliser (kg N):

N

To wheat	To barley	Wheat	Barley
63	50	63	50
126	100	126	100
189	150	189	150
252	200	252	200

NOTE: Ley and potatoes receive standard N only, residues of dressings to cereals are tested (NRESID).

75/W/RN/13

Basal applications: All crops: P2O5 at 130 kg, K2O at 260 kg as (0:14:28), half ploughed in, half applied to the plough furrow.

Standard applications:

Leys: N at 60 kg, as 'Nitro-Chalk', in seedbed and repeated after sowing.

Weedkiller: Amfotriazole at 4.5 kg in 280 l.

Potatoes: N at 150 kg as 'Nitro-Chalk'. Weedkiller: Linuron at 1.2 kg plus paraquat at 0.28 kg ion in 280 l. Insecticide: Demeton-s-methyl at 0.25 kg in 280 l. Fungicide: Mancozeb at 1.3 kg in 390 l.

Wheat: Weedkillers: Aminotriazole at 4.5 kg in 280 l. Ioxynil at 0.63 kg with mecoprop at 1.9 kg in 280 l.

Barley: Weedkillers: Aminotriazole at 4.5 kg in 280 l. Ioxynil at 0.52 kg with mecoprop at 1.6 kg in 280 l.

Seed: Leys: Italian ryegrass sown at 40 kg.

Potatoes: Pentland Crown.

Wheat: Cappelle sown at 200 kg.

Barley: Julia, dressed with ethirimol, sown at 160 kg.

Cultivations, etc.: - All plots: Half PK applied, ploughed: 30 Oct, 1974.

Remaining PK applied: 8 Nov. Spring-tine cultivated: 27 Nov.

Leys: Aminotriazole applied: 11 Sept., 1974. Spring-tine cultivated four times with crumbler on the second and fourth occasions: 26 Feb, 3 Mar, 24 Apr, 28 Apr, 1975. N applied: 18 Apr, 11 June. Seed sown: 2 May. Topped: 8 July, 21 Aug.

Potatoes: Spring-tine cultivated three times on the second occasion with crumbler: 26 Feb, 3 Mar, 24 Apr, 1975. N applied: 18 Apr. Deep-tine cultivated: 25 Apr. Rotary cultivated, potatoes planted: 5 May. Linuron with paraquat applied: 22 May. Grubbed: 23 June. Rotary ridged: 24 June. Insecticide applied: 25 June. Fungicide applied: 15 July. Haulm mechanically destroyed: 26 Sept. Sprayed with undiluted B.O.V. at 160 l: 2 Oct. Lifted: 6 Oct.

Wheat: Seed sown: 27 Nov, 1974. N applied: 24 Mar, 1975. Weedkiller applied: 8 May. Combine harvested: 12 Aug.

Barley: Spring-tine cultivated twice, second time with crumbler: 26 Feb, 3 Mar, 1975. Seed sown: 4 Mar. N applied: 19 Mar. Spring-tine cultivated with crumbler three times, seed resown: 1 May. Weedkiller applied: 4 June. Combine harvested: 18 Aug.

- NOTES: (1) No cuts were taken from the leys because of poor growth in a dry year.  
(2) The barley was resown because of poor germination and bird damage on the first sowing.  
(3) Estimates of eyespot (*Cercospora herpotrichoides*) and take-all (*Gaeumannomyces graminis*) were made on both cereal crops.



75/W/RN/13

POTATOES

WHEAT SITE

TOTAL TUBERS TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

NRES ID	63	126	189	252	MEAN
	25.4	23.9	24.2	20.3	23.4

PERCENTAGE WARE 3.81CM (1.5 INCH) RIDDLE

\*\*\* TABLES OF MEANS \*\*\*

NRES ID	63	126	189	252	MEAN
	97.4	95.7	96.3	95.6	96.2

BARLEY SITE

TOTAL TUBERS TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

NRES ID	50	100	150	200	MEAN
	29.5	29.8	26.6	25.1	27.8

PERCENTAGE WARE 3.81CM (1.5 INCH) RIDDLE

\*\*\* TABLES OF MEANS \*\*\*

NRES ID	50	100	150	200	MEAN
	97.5	97.4	97.5	97.4	97.5

PLOT AREA HARVESTED 0.00139

75/W/RN/13

WINTER WHEAT

GRAIN TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

	N	63	126	189	252	MEAN
PREVCROP						
C/C/L/P		2.75	3.36	3.45	3.43	3.25
C/L/P/C		1.46	2.97	3.39	2.83	2.66
L/P/C/C		1.91	3.15	3.14	2.80	2.75
C/C/C/C		1.81	3.04	3.22	2.77	2.71
MEAN		1.98	3.13	3.30	2.96	2.84

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	N	PREVCROP*
		N
SED	0.150	0.300

\* WITHIN THE SAME LEVEL OF PREVCROP ONLY

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	3	0.153	5.4
BLOCK.WP.SP	12	0.300	10.6

GRAIN MEAN DM% 88.2

STRAW TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

	N	63	126	189	252	MEAN
PREVCROP						
C/C/L/P		2.41	4.00	4.52	4.89	3.95
C/L/P/C		1.25	2.85	3.84	3.57	2.88
L/P/C/C		1.86	2.50	3.51	3.74	2.90
C/C/C/C		2.92	3.10	3.62	3.24	3.22
MEAN		2.11	3.11	3.87	3.86	3.24

STRAW MEAN DM% 89.8

SUB PLOT AREA HARVESTED 0.00277

75/W/RN/13

BARLEY

GRAIN TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

	N	50	100	150	200	MEAN
PREVCROP						
C/C/L/P		2.08	3.13	3.23	2.88	2.83
C/L/P/C		1.73	2.83	2.94	2.82	2.58
L/P/C/C		1.84	2.47	2.73	2.87	2.49
C/C/C/C		1.64	2.11	2.14	2.29	2.04
MEAN		1.82	2.64	2.77	2.71	2.48

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	N	PREVCROP*
		N
SED	0.117	0.234

\* WITHIN THE SAME LEVEL OF PREVCROP ONLY

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	3	0.286	11.5
BLOCK.WP.SP	12	0.234	9.4

GRAIN MEAN DM% 86.9

STRAW TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

	N	50	100	150	200	MEAN
PREVCROP						
C/C/L/P		1.65	2.57	2.57	2.41	2.30
C/L/P/C		1.41	2.42	2.41	2.58	2.21
L/P/C/C		1.75	2.18	2.35	2.63	2.23
C/C/C/C		1.26	1.76	1.96	1.98	1.74
MEAN		1.52	2.23	2.32	2.40	2.12

STRAW MEAN DM% 81.6

SUB PLOT AREA HARVESTED 0.00277