
Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1975



Full Table of Content

75/ R/CS/41 - Cultivations & Soil Invertebrates - Old Grass, New Grass

Rothamsted Research

Rothamsted Research (1976) 75/ R/CS/41 - Cultivations & Soil Invertebrates - Old Grass, New Grass; Yields Of The Field Experiments 1975, pp 187 - 189 - **DOI**:

https://doi.org/10.23637/ERADOC-1-141

75/R/CS/41

CULTIVATIONS AND SOIL INVERTEBRATES

Object: To study the effects of cultivations on yields of grass and on populations of soil animals - Road Piece.

Sponsor: C.A. Edwards.

The seventh year, old grass, new grass.

For previous years see 69/R/CS/41(t), 70/R/CS/41(t) and 71-74/R/CS/41.

Design: 4 blocks of 8 plots randomisation restricted.

Whole plot dimensions: 6.40 x 7.32.

| Treatments: Cultivations and reseeding: | CULTIVIN |
|---|----------|
| No treatments to old grass (two plots per block) | 0 |
| Grass ploughed up:- | |
| In spring 1969, reseeded after fewest | |
| cultivations needed to produce a seedbed | SF |
| In spring 1969, reseeded after many seedbed | |
| cultivations | SM |
| In autumn 1969, reseeded spring 1970 after | ONI |
| | |
| many seedbed cultivations | AM |
| Every spring since 1969, reseeded each year | |
| after fewest cultivations needed to | |
| produce a seedbed | SFR |
| Every spring since 1969, reseeded each year | 5111 |
| after many seedbed cultivations | CATE: |
| Every autumn since 1969, reseeded every following | SMR |
| | |
| spring after many seedbed cultivations | AMR |

Seeds mixture for 1975: 6.7 kg \$215 Meadow Fescue, 4.5 kg Contessa Meadow Fescue, 4.5 kg \$48 Timothy, 1.7 kg Nina certified Wild White Clover, 0.6 kg Old Pasture Wild White Clover. Mixture sown at 24 kg.

Basal applications: Manures: (0:14:28) at 500 kg in winter, (25:0:16) at 440 kg in spring, (25:0:16) at 220 kg after each cut except the last.

Cultivations, etc.:- Basal PK applied: 15 Jan, 1975. AMR plots ploughed: 24 Jan. Basal NK applied: 5 Mar. SMR and SFR plots ploughed: 19 Mar. Disced AMR and SMR plots six times, SFR plots three times: 5 May. Power harrowed all plots to be sown: 6 May. Seed sown: 9 May. AMR, SMR, SFR plots topped: 14 July. Other plots cut three times: 28 May, 31 July, 7 Nov. NK applied to all plots except AMR, SMR, SFR: 16 June, and to all plots: 18 Aug.

NOTE: Soil cores were taken to assess total fauna on 7 May, 16 Oct, and quadrats were sampled on each plot for earthworms on 29 May, 16 Oct.

75 /R /CS /41

1ST CUT (28/5/75) DRY MATTER TONNES/HECTARE

*** TABLES OF MEANS ***

CULTIVIN 0 SF SM AM MEAN 4.67 4.45 4.19 4.75 4.55

*** STANDARD ERRORS OF DIFFERENCES OF MEANS ***

TABLE CULTIVTN

SED 0.231(1)
0.267(2)

- (1) O V ANY REMAINDER
- (2) ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

S TRATUM DF SE CV%

BLCCK.WP 13 0.378 8.3

1ST CUT MEAN DM% 18.3

2ND CUT (31/7/75) DRY MATTER TONNES/HECTARE

*** TABLES OF MEANS ***

CULTIVIN 0 SF SM AM MEAN 0.99 1.52 2.04 1.90 1.49

*** STANDARD ERRORS OF DIFFERENCES OF MEANS ***

TABLE CULTIVTN

SED 0.254(1)
0.293(2)

- (1) O V ANY REMAINDER
- (2) ANY OF REMAINDER

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

 STRATUM
 DF
 SE
 CV%

 BLCCK.WP
 13
 0.414
 27.9

2ND CUT MEAN DM% 31.2

188

75/R/CS/41

3RD CUT (7/11/75) DRY MATTER TONNES/HECTARE

*** TABLES OF MEANS ***

CULTIVIN 0 SF SM AM MEAN 0.73 0.78 0.79 0.69 0.74

*** STANDARD ERRORS OF DIFFERENCES OF MEANS ***

TABLE CULTIVTN

SED 0.116(1)
0.134(2)

(1) O V ANY REMAINDER

(2) ANY OF REMAINDER

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV%

BLCCK.WP 13 0.190 25.6

3RD CUT MEAN DM% 24.2

TOTAL OF 3 CUTS DRY MATTER TONNES/HECTARE

*** TABLES OF MEANS ***

CULTIVTN 0 SF SM AM MEAN 6.39 6.74 7.01 7.3 6.73

*** STANDARD ERRORS OF DIFFERENCES OF MEANS ***

TABLE CULTIVTN

SED 0.410(1)
0.473(2)

(1) O V ANY OF REMAINDER

(2) ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV%

BLCK.WP 13 0.669 9.9

TOTAL OF 3 CUTS MEAN DM% 24.5

PLOT AREA HARVESTED 0.00074

189