


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1975

[Full Table of Content](#)



75/ W/CS/99 - Effects of Breaks on Take-all - Barley, Beans

Rothamsted Research

Rothamsted Research (1976) 75/ W/CS/99 - *Effects of Breaks on Take-all - Barley, Beans* ; Yields Of The Field Experiments 1975, pp 195 - 196 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/W/CS/99

EFFECTS OF BREAKS ON TAKE-ALL

Object: To study the phenomenon of 'take-all' (*Gaeumannomyces graminis*) decline in barley - Woburn Butt Furlong.

Sponsor: D. Hornby.

The fourth year, barley, spring beans.

For previous years see 72/W/CS/99(t) and 73-74/W/CS/99.

Design: 2 blocks of 9 plots.

Whole plot dimensions: 5.33 x 15.2.

Treatments: Previous crops:-

PREVCROP

1968-71	1972	1973	1974		
B	B	B	B	(4 plots per block)	B/B/B/B
B	F	B	B		B/F/B/B
B	F	BE	B		B/F/BE/B
B	B	F	BE		B/B/F/BE

B = Barley, BE = Beans, F = Fallow

Standard applications:

Barley: Manures: (20:14:14) at 420 kg combine drilled. Weedkiller: Ioxynil at 0.53 kg and mecoprop at 1.6 kg in 280 l.

Spring beans: Manures: (0:14:28) at 400 kg placement drilled. Weedkiller: Simazine at 0.84 kg in 280 l. Insecticide: Demeton-s-methyl at 0.20 kg in 390 l.

Seed: Barley: Julia, dressed with ethirimol, sown at 160 kg.
Spring beans: Minden, sown at 220 kg.

Cultivations, etc.:-

All plots: Ploughed: 11-13 Dec, 1974. Spring-tine cultivated: 25 Feb, 1975.

Barley: Seed sown: 28 Feb, 1975. Weedkiller applied: 22 May.
Combine harvested: 12 Aug.

Spring beans: Spring-tine cultivated with crumbler: 21 Apr, 1975. Seed sown: 23 Apr. Weedkiller applied: 26 Apr. Insecticide applied: 14 July. Combine harvested: 26 Aug.

Fallow: Spring-tine cultivated with crumbler: 21 Apr, 1975. Spring-tine cultivated: 22 May.

NOTE: Soil samples were taken before sowing and after harvest and plant samples in July for incidence of 'take-all' (*Gaeumannomyces graminis*).

75/W/CS/99

*** TABLES OF MEANS ***

BARLEY GRAIN TONNES/HECTARE

PREVCROP	B/B/B/B	B/F/B/B	B/F/BE/B	B/B/F/BE	MEAN
	1.42	2.08	2.16	2.19	1.73

*** STANDARD ERRORS OF DIFFERENCES OF MEANS ***

TABLE	PREVCROP	
SED	0.270	B/B/B/B V ANY OF REMAINDER
	0.341	ANY OF REMAINDER

--- STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ***

STRATUM	DF	SE	CV%
BLOCK.WP	9	0.341	19.7

GRAIN MEAN DM% 87.0

PLOT AREA HARVESTED 0.00434

SPRING BEANS GRAIN TONNES/HECTARE

MEAN 0.76

MEAN DM% 86.7