

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1975

[Full Table of Content](#)



## 75/ R/CS/109 - Benomyl & Sclerotinia - Winter Beans

### Rothamsted Research

Rothamsted Research (1976) 75/ R/CS/109 - Benomyl & Sclerotinia - Winter Beans ; Yields Of The Field Experiments 1975, pp 201 - 202 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/CS/109

BENOMYL AND SCLEROTINIA

Object: To study the effects of three fungicides on yield and incidence of Sclerotinia rot in winter beans - Fosters O & E III.

Sponsor: J.F. Jenkyn.

The third year, winter beans.

For previous years see 73-74/R/CS/109.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 4.06 x 4.27.

Treatments: Fungicide:	FUNGICIDE
None (3 plots per block)	O
Benomyl, two sprays each to 1973 and 1974 crops, one in 1975 (3 plots per block)	B(2 2) 1
Benomyl, five sprays each to 1973 and 1974 crops, two in 1975	B(5 5) 2
Benomyl + RP 26019, no sprays in 1973 and 1974, two in 1975	BR(O) 2
RP 26019, no sprays in 1973 and 1974, two in 1975	R(O) 2
Captafol, none in 1973, five sprays in 1974, two in 1975	C(O 5) 2

- NOTES: (1) Crop in 1973 and 1974 was red clover, infected with Sclerotinia.  
(2) Sprays were applied on 7 Feb and 9 May, 1975 (B(2 2) 1 7 Feb only).  
(3) Rates of application, in 340 l:-  
Benomyl at 0.56 kg. RP 26019 at 0.56 kg a.i. and captafol at 1.34 kg.

Basal applications: Manures: (0:14:28) at 850 kg.

Seed: Throws MS, sown at 380 kg.

Cultivations, etc.: - Clover ploughed: 11 July, 1974. Rotary cultivated twice: 30 July and twice: 20 Aug. Spring-tine cultivated: 28 Aug. Ploughed: 16 Sept. PK fertiliser applied: 28 Oct. Power harrowed: 29 Oct. Seed sown: 4 Nov. Combine harvested: 6 Aug, 1975.

NOTE: The crop was sown late and very few plants became infected with Sclerotinia.

75/R/CS/109

\*\*\* TABLES OF MEANS \*\*\*

GRAIN TONNES/HECTARE

FUNGCIDE	0	B(2 2) 1	B(5 5) 2	BR(0) 2	R(0) 2	C(0 5) 2	MEAN
	2.94	3.10	3.01	3.07	3.25	3.00	3.04

\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*

TABLE	FUNGCIDE	
SED	0.157	0 v B(2 2) 1
SED	0.223	0 or B(2 2) 1 v ANY OF REMAINDER
SED	0.273	ANY OF REMAINDER

\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	22	0.334	11.0

GRAIN MEAN DM% 87.5

PLOT AREA HARVESTED 0.00065