
Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1975



Full Table of Content

75/ R/CS/123 - Effects of Viruses - Ryegrass, Clover

Rothamsted Research

Rothamsted Research (1976) 75/ R/CS/123 - Effects of Viruses - Ryegrass, Clover; Yields Of The Field Experiments 1975, pp 209 - 212 - DOI: https://doi.org/10.23637/ERADOC-1-141

75/R/CS/123

EFFECTS OF VIRUSES

Object: To study the effects of virus infection on yield and persistence of ryegrass and red clover in pure and mixed stands - Garden Plot 14.

Sponsors: A.J. Cockbain, R.T. Plumb.

The third year, red clover and Italian ryegrass.

For previous years see 73-74/R/CS/123.

Design: 4 randomised blocks of 8 plots split into 2.

Whole plot dimensions: 3.35 x 1.22.

Treatments: All combinations of:-

Whole plots: 1. Crop and inoculation with virus in 1973: CROP VIR

Red clover uninoculated Italian ryegrass uninoculated	CLOVER/O GRASS/O
Red clover and Italian ryegrass (mixed within rows) uninoculated	CLOGR/O
Red clover inoculated with Pea Mosaic Virus (PMV)	CLOVER/P
Italian ryegrass inoculated with Ryegrass Mosaic Virus (RMV)	GRASS/R
Red clover and Italian ryegrass (mixed) inoculated with PMV	CLOGR/P
Red clover and Italian ryegrass (mixed) inoculated	THE RESIDENCE OF THE
with RMV	CLOGR/R
Red clover and Italian ryegrass (mixed) inoculated	

NOTE: Inoculation with PMV was unsuccessful.

with PMV and RMV

Sub plots:	2. Aldicarb (kg) after each cut:	ALDICARB
	None	0

10 10

CLOGR/PR

Basal applications: Manures: (25:0:16) at 270 kg to grass plots.

Cultivations, etc.:- NK applied: 19 June, 1975. Cut twice: 9 June, 25 July.

NOTES: (1) Mites (Abacarus hystrix) were counted on 28 Apr and 23 July on grass plots.

(2) Grass was scored for Ryegrass Mosaic Virus on 20 May.

75 /R /CS /123

1ST CUT (6/6/75) DRY MATTER TONNES/HECTARE

*** TABLES OF MEANS ***

ALDICARB	0	10	MEAN
CROP VIR			
CLOVER /O	8.37	8.45	8.41
GRASS/0	8.25	7.63	7.94
CLOGR/O	9.53	8.77	9.15
CLOVER /P	8.35	8.31	8.33
GRASS/R	8.08	7.36	7.72
CLOGR/P	9.40	8.76	9.08
CLOGR/R	8.84	8.93	8.89
CLOGR/PR	8.92	8.57	8.75
MEAN	8.72	8.35	8.53

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	CRO	P VIR	AL	DICARI	B CROP	
REP	 	8		32	2	4
SED		0.440		0.160	0	.544
EXCEP T CROP	COMPARING	MEANS	WITH	SAME		OF:

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM	DF	SE	CV%
BLCCK.WP	21	0.623	7.3
BLCCK.WP.SP	24	0.638	7.5

1ST CUT MEAN DM% 25.2

75/R/CS/123

2ND CUT (25/7/75) DRY MATTER TONNES/HECTARE

*** TABLES OF MEANS ***

ALDICARB	0	10	MEAN
CROP VIR			
CLOVER/O	3.14	3.89	3.51
GRASS /O	1.89	1.46	1.68
CLOGR/O	3.04	3.09	3.06
CLOVER/P	3.27	3.76	3.51
GRASS/R	1.57	1.65	1.61
CLGR/P	2.96	2.92	2.94
CLOGR/R	3.30	3.10	3.20
CLOGR/PR	2.76	3.27	3.01
MEAN	2.74	2.89	2.82

***** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	CRO	P VIR	AL	DICARE	CROP VIR ALDICARB
REP SED EXCEPT CROP		8 0.378 MEANS	WITH	32 0.122 SAME	

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM	DF	SE	CV%
BLCCK.WP	21	0.535	19.0
BLCCK.WP.SP	24	0.486	17.3

2ND CUT MEAN DM% 26.8

75/R/CS/123

TOTAL OF 2 CUTS DRY MATTER TONNES/HECTARE

*** TABLES OF MEANS ***

ALDICARB	0	10	MEAN
CROP VIR			
CLOVER/O	11.51	12.34	11.92
GRASS/O	10.14	9.10	9.62
CLOGR/O	12.56	11.86	12.21
CLOVER /P	11.63	12.06	11.84
GRASS/R	9.65	9.01	9.33
CLOGR/P	12.36	11.68	12.02
CLOGR/R	12.14	12.03	12.09
CLOGR/PR	11.68	11.84	11.76
MEAN	11.46	11.24	11.35

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE		CRO	P VIR	AL	DICARI		VIR
REP			8		32	2	4
SED			0.695		0.248	8 0	. 854
EXCEPT	WHEN	COMPARING	MEANS	WITH	SAME	LEVEL(S)	OF:
CROP	VIR					0	.702

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM	DF	SE	CV%
BLCCK.WP	21	0.982	8.7
BLCCK.WP.SP	24	0.993	8.8

TOTAL OF 2 CUTS MEAN DM% 26.0

PLOT AREA HARVESTED 0.00011