

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1975

[Full Table of Content](#)



75/ R/CS/135 - Direct Drilling & Diseases - Winter Barley

Rothamsted Research

Rothamsted Research (1976) 75/ R/CS/135 - *Direct Drilling & Diseases - Winter Barley* ; Yields Of The Field Experiments 1975, pp 224 - 225 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/CS/135

DIRECT DRILLING AND DISEASES

Object: To study the effects of direct drilling on pathogens and yield of winter barley - Meadow.

Sponsor: R.D. Prew.

The second year, winter barley.

For previous year see 74/R/CS/135.

Design: 2 randomised blocks of 3 plots.

Whole plot dimensions: 26.7 x 30.5.

Treatments: Cultivations:-	CULTIVIN
Ploughed, spring-tine cultivated, drilled (P)	PLOUGH
Paraquat to stubble, chisel ploughed, rotary harrowed, drilled (C)	CHISEL
Paraquat to stubble, direct drilled (D)	DIRECT

NOTE: Paraquat was applied at 0.84 kg ion in 220 l.

Basal applications: Manures: (10:24:24) at 250 kg combine drilled. 'Nitro-Chalk' in spring at 375 kg. Weedkiller: Dicamba with mecoprop and MCPA ('Banlene Plus' at 5.6 l in 220 l).

Seed: Maris Otter seed dressed fungicide only, sown at 160 kg (P & C), 190 kg (D).

Cultivations, etc.:- Paraquat applied to C and D plots: 14 Oct, 1974.
P plots ploughed: 15 Oct. C plots chisel ploughed: 17 Oct. D plots direct drilled: 24 Oct. P plots spring-tine cultivated, C plots rotary harrowed and seed sown on P and C plots: 25 Oct. N applied: 21 Apr, 1975. Weedkiller applied: 12 May. Combine harvested: 30 July. Previous crops: Barley 1973 and 1974.

NOTES: (1) Much straw trash was left on the surface before drilling, to aid study of foliar diseases. Because of this the crop established badly on D plots and became infested with grass weeds.
(2) Root and foliar diseases were assessed.

75/R/CS/135

*** TABLES OF MEANS ***

GRAIN TONNES/HECTARE

CULTIVTN	PLOUGH	CHISEL	DIRECT	MEAN
	4.01	4.38	3.08	3.83

GRAIN MEAN DM% 87.5

PLOT AREA HARVESTED 0.00929