


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1975

[Full Table of Content](#)



75/ R/CS/140 - Chemical Reference Plots - Barley

Rothamsted Research

Rothamsted Research (1976) 75/ R/CS/140 - *Chemical Reference Plots - Barley* ; Yields Of The Field Experiments 1975, pp 231 - 234 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of four agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of continuous barley - Long Hoos V 3.

Sponsors: G.G. Briggs, N. Walker, R. MacDonald.

The second year, barley.

For previous year see 74/R/CS/140.

Design: Two replicates of 2 x 2 x 2 x 2 fully randomised.

Whole plot dimensions: 4.06 x 4.57.

Treatments, applied cumulatively to 1974 except WEEDKILLR - 1974 only:

All combinations of:-

1. Fungicide:

FUNGCIDE

None

NONE

Benomyl at 4 kg applied 22 Apr, 1975

BENOMYL

2. Insecticide:

INSCTCDE

None

NONE

Chlorfenvinphos at 2 kg applied 22 Apr, 1975

CHLORFEN

3. Nematicide:

NEMACIDE

None

NONE

Aldicarb at 6 kg applied 22 Apr, 1975

ALDICARB

4. Weedkiller:

WEEDKILLR(74)

None

NONE

Chlortoluron at 2 kg applied 1974 only

CHLORTOL

Basal applications: Manures: (20:14:14) at 500 kg. Weedkillers: Dicamba with mecoprop and MCPA ('Tetralex Plus' at 7.0 l in 340 l).

Seed: Julia - undressed, sown at 160 kg.

Cultivations, etc.: Ploughed: 19 Sept, 1974. Power harrowed, NPK applied, seed sown: 22 Apr, 1975. Weedkiller applied: 6 June. Combine harvested: 13 Aug.

- NOTES: (1) Mildew and aphid numbers were assessed in May, June and July, and barley yellow dwarf virus in July. Chemical residues in the soil were assessed throughout the season and aldicarb in the crop in July.
(2) There was evidence of a linear trend across the site, and yields adjusted for this trend are presented.

75/R/CS/140

GRAIN TONNES/HECTARE

*** TABLES OF MEANS ***

| INSCTCDE FUNGCIDE | NONE | CHLORFEN | MEAN |
|--------------------------|------|----------|------|
| NONE | 5.01 | 5.08 | 5.05 |
| BENOMYL | 5.22 | 5.39 | 5.31 |
| MEAN | 5.12 | 5.23 | 5.18 |
| NEMACIDE FUNGCIDE | NONE | ALDICARB | MEAN |
| NONE | 4.75 | 5.34 | 5.05 |
| BENOMYL | 4.96 | 5.66 | 5.31 |
| MEAN | 4.86 | 5.50 | 5.18 |
| NEMACIDE INSCTCDE | NONE | ALDICARB | MEAN |
| NONE | 4.88 | 5.35 | 5.12 |
| CHLORFEN | 4.83 | 5.64 | 5.23 |
| MEAN | 4.86 | 5.50 | 5.18 |
| WEEDKLIR(74) FUNGCIDE | NONE | CHLORTOL | MEAN |
| NONE | 5.05 | 5.05 | 5.05 |
| BENOMYL | 5.13 | 5.49 | 5.31 |
| MEAN | 5.09 | 5.27 | 5.18 |
| WEEDKLIR(74) INSCTCDE | NONE | CHLORTOL | MEAN |
| NONE | 5.07 | 5.17 | 5.12 |
| CHLORFEN | 5.10 | 5.37 | 5.23 |
| MEAN | 5.09 | 5.27 | 5.18 |
| WEEDKLIR(74) NEMACIDE | NONE | CHLORTOL | MEAN |
| NONE | 4.77 | 4.94 | 4.86 |
| ALDICARB | 5.40 | 5.60 | 5.50 |
| MEAN | 5.09 | 5.27 | 5.18 |

75/R/CS/140

GRAIN TONNES/HECTARE

| | | | | |
|--------------------|------|----------|---------------|------|
| INSCTCDE | NONE | CHLORFEN | | |
| NEMACIDE | NONE | ALDICARB | NONE ALDICARB | |
| FUNGicide | | | | |
| NONE | 4.70 | 5.33 | 4.81 | 5.35 |
| BENOMYL | 5.06 | 5.38 | 4.85 | 5.93 |
| INSCTCDE | NONE | CHLORFEN | | |
| WEEDKLLR(74) | NONE | CHLORTOL | NONE CHLORTOL | |
| FUNGicide | | | | |
| NONE | 5.04 | 4.99 | 5.05 | 5.10 |
| BENOMYL | 5.10 | 5.35 | 5.15 | 5.63 |
| NEMACIDE | NONE | ALDICARB | | |
| WEEDKLLR(74) | NONE | CHLORTOL | NONE CHLORTOL | |
| FUNGicide | | | | |
| NONE | 4.72 | 4.78 | 5.37 | 5.31 |
| BENOMYL | 4.82 | 5.09 | 5.43 | 5.89 |
| NEMACIDE | NONE | ALDICARB | | |
| WEEDKLLR(74) | NONE | CHLORTOL | NONE CHLORTOL | |
| INSCTCDE | | | | |
| NONE | 4.86 | 4.91 | 5.28 | 5.43 |
| CHLORFEN | 4.69 | 4.97 | 5.52 | 5.76 |
| NEMACIDE | NONE | ALDICARB | | |
| WEEDKLLR(74) | NONE | CHLORTOL | NONE CHLORTOL | |
| FUNGicide INSCTCDE | | | | |
| NONE | NONE | 4.75 | 4.65 | 5.32 |
| CHLORFEN | 4.69 | 4.92 | 5.41 | 5.28 |
| BENOMYL | NONE | 4.96 | 5.16 | 5.23 |
| CHLORFEN | 4.69 | 5.02 | 5.62 | 6.25 |

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

| TABLE | FUNGicide | INSCTCDE | NEMACIDE | WEEDKLLR(74) |
|-------|---------------------------------------|--------------------------------------|-----------------------------------|---------------------------------------|
| SED | 0.129 | 0.135 | 0.128 | 0.133 |
| TABLE | FUNGicide INSCTCDE | FUNGicide NEMACIDE | INSCTCDE NEMACIDE | FUNGicide WEEDKLLR(74) |
| SED | 0.185 | 0.182 | 0.189 | 0.186 |
| TABLE | INSCTCDE WEEDKLLR(74) | NEMACIDE WEEDKLLR(74) | FUNGicide INSCTCDE NEMACIDE | FUNGicide INSCTCDE WEEDKLLR(74) |
| SED | 0.191 | 0.188 | 0.261 | 0.265 |
| TABLE | FUNGicide NEMACIDE WEEDKLLR(74) | INSCTCDE NEMACIDE WEEDKLLR(74) | FUNGicide INSCTCDE NEMACIDE | FUNGicide INSCTCDE WEEDKLLR(74) |
| SED | 0.261 | 0.268 | 0.372 | |

75/R/CS/140

GRAIN TONNES/HECTARE

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

| STRATUM | DF | SE | CV% |
|---------|----|-------|-----|
| WP | 15 | 0.362 | 7.0 |

GRAIN MEAN DM% 89.1

PLOT AREA HARVESTED 0.00075