


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1975

[Full Table of Content](#)



75/ R/CS/155 - Rainfall & Leaching - Spring Wheat

Rothamsted Research

Rothamsted Research (1976) 75/ R/CS/155 - *Rainfall & Leaching - Spring Wheat* ; Yields Of The Field Experiments 1975, pp 251 - 252 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/CS/155

RAINFALL AND LEACHING

Object: To study the leaching of nitrate nitrogen by a range of winter rainfall rates and to relate leaching and responses to fertiliser nitrogen applied to spring wheat - Long Hoos VI/VII 4.

Sponsors: T.M. Addiscott, D. Cox.

The first year, spring wheat.

Design: 2 blocks of 16 plots, randomisation restricted.

Whole plot dimensions: 2.13 x 2.13.

Treatments: All combinations of:

1. 'Rainfall' (mm), October-February inclusive: RAINFALL

177	177
258	258
396	396
484	484

2. Nitrogen fertiliser, applied 16 May (kg N):

	N
None	0
40	40
80	80
120	120

NOTE: Natural rainfall in the period October-February was 396 mm. Smaller rates were achieved by covering and the larger rate by supplementing natural rainfall with irrigation.

Basal applications: Manures: (0:20:20) at 250 kg combine drilled.

Weedkiller: Dicamba with mecoprop and MCPA ('Tetralix Plus' at 7.0 l in 340 l).

Seed: Kleiber, sown at 190 kg.

Cultivations, etc.: Ploughed: 19 Sept, 1974. Power harrowed: 20 Sept.

Hand dug: 26 Mar, 1975. Power harrowed and seed sown: 22 Apr.

Weedkiller applied: 6 June. Combine harvested: 28 Sept. Previous crops: Fallow 1973 and 1974.

NOTES: (1) Concentrations of nitrate and ammonium ions at two soil depths were determined at monthly intervals, and nitrate concentration at additional levels on 11 March.

(2) N content of grain was determined.

(3) There was evidence of a fertility trend across the site and yields adjusted for this trend are presented.

75/R/CS/155

GRAIN TONNES/HECTARE

*** TABLES OF MEANS ***

	N	0	40	80	120	MEAN
RAINFALL						
177		1.51	2.45	2.35	2.68	2.25
258		1.67	2.42	2.79	2.28	2.29
396		1.73	2.03	2.10	2.10	1.99
484		1.70	1.90	2.47	2.13	2.05
MEAN		1.65	2.20	2.42	2.30	2.14

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	RAINFALL	N	RAINFALL N
SED	0.168	0.169	0.330

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	14	0.317	14.8

GRAIN MEAN DM% 87.6

PLOT AREA HARVESTED 0.00035