


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1975

[Full Table of Content](#)



75/ S/CS/1 - Varieties, N & Ccc - Wheat

Rothamsted Research

Rothamsted Research (1976) 75/ S/CS/1 - Varieties, N & Ccc - Wheat ; Yields Of The Field Experiments 1975, pp 279 - 282 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/S/CS/1

VARIETIES. N AND CCC

Object: To study the effects of nitrogen fertiliser, at a range of rates and times, and chlormequat (CCC) on the yield of two varieties of winter wheat - Saxmundham, Oldershaw's and Garner's plots.

Sponsors: F.V. Widdowson, A.E. Johnston.

The tenth year, winter wheat.

For previous years see 66/C/30(t), 67/C/23(t), 68/C/39, 69-70/S/CS/1, 71/S/CS/1(t), 72/S/CS/1(t) and 73-74/S/CS/1.

Design: A single replicate of 4 x 2 x 2 x 2 in 4 blocks of 4 plots, each split lengthways into 2, plus one additional plot per block. Additionally all the plots are split breadthways into 3.

Whole plot dimensions: 5.49 x 40.2.

Treatments: All combinations of:-

Whole plots (All sown at a seed rate of 170 kg with 13 cm (5 inches)

between the rows): 1. Number of previous continuous wheat crops:

6	6 WHEAT
7	7 WHEAT
8	8 WHEAT
9	9 WHEAT

2. Chlormequat (kg):

None	0.0
1.7 in 340 l	1.7

Half plots:

3. Times of applying nitrogen fertiliser: N TIME

Single dressing (5 May)	SINGLE
Divided dressing (Half 23 April, half 15 May)	DIVIDED

4. Varieties: VARIETY

Cappelle	CAPPELLE
Maris Huntsman	HUNTSMAN

75/S/CS/1

Pairs of sixth plots: 5. Rates of nitrogen fertiliser in addition to 62 kg N in autumn (28 Oct) (kg N): N RATE

50	50
100	100
150	150

Together with one extra plot per block which had 5 previous wheat crops and was sown with Cappelle at a seed rate of 180 kg with 20 cm (8 inches) between the rows and tested all combinations of: EXTRA

Half plots: 1. Time of applying nitrogen fertiliser: N TIME

23 Apr	APRIL
15 May	MAY

Pairs of sixth plots: 2. Rate of nitrogen fertiliser (kg N): N RATE

50	50
100	100
150	150

NOTE: 62 kg N, 31 kg P₂O₅, 31 kg K₂O was broadcast at drilling as (20:10:10) to all plots except EXTRA. EXTRA plots were to have tested autumn N as in 74/S/CS/1 but bad weather prevented this. A test of early v late spring N was substituted and these plots received a balancing dressing of 31 kg P₂O₅ and 31 kg K₂O as (0:20:20).

Basal applications: Manures: (0:20:20) at 1260 kg applied to stubble before ploughing. Weedkillers: Ioxynil at 0.63 kg with mecoprop at 1.9 kg in 450 l.

Cultivations, etc.: - Basal PK applied: 10 Sept, 1974. Ploughed: 16 Sept. NPK applied to all plots except EXTRA and all seed sown: 28 Oct. Balancing PK applied to EXTRA plots: 22 Apr, 1975. Weedkillers applied: 30 Apr. Chlormequat applied: 15 May. Combine harvested: 19 Aug.

75/5/CS/1

GRAIN TONNES/HECTARE

*** TABLES OF MEANS ***

PREVCROP	6 WHEAT	7 WHEAT	8 WHEAT	9 WHEAT	MEAN
CCC					
0.0	3.81	3.99	3.76	3.84	3.85
1.7	4.35	4.05	4.21	3.75	4.09
MEAN	4.08	4.02	3.99	3.80	3.97

PREVCROP	6 WHEAT	7 WHEAT	8 WHEAT	9 WHEAT	MEAN
N TIME					
SINGLE	3.99	3.62	3.83	3.65	3.77
DIVIDED	4.17	4.42	4.14	3.95	4.17
MEAN	4.08	4.02	3.99	3.80	3.97

PREVCROP	6 WHEAT	7 WHEAT	8 WHEAT	9 WHEAT	MEAN
VARIETY					
CAPPELLE	4.12	4.11	4.02	3.67	3.98
HUNTSMAN	4.04	3.94	3.95	3.93	3.97
MEAN	4.08	4.02	3.99	3.80	3.97

PREVCROP	6 WHEAT	7 WHEAT	8 WHEAT	9 WHEAT	MEAN
N RATE					
50	2.98	3.04	2.69	2.76	2.87
100	3.81	4.24	4.08	3.93	4.02
150	5.45	4.78	5.19	4.70	5.03
MEAN	4.08	4.02	3.99	3.80	3.97

GRAIN MEAN DM% 85.1

PLOT AREA HARVESTED 0.00355

75/5/CS/1

EXTRA

GRAIN TONNES/HECTARE

*** TABLES OF MEANS ***

N RATE	50	100	150	MEAN
N TIME				
APRIL	2.80	3.62	4.62	3.68
MAY	2.84	3.62	4.49	3.65
MEAN	2.82	3.62	4.55	3.67

GRAIN MEAN DM% 85.0

STRAW TONNES/HECTARE

*** TABLES OF MEANS ***

N RATE	50	100	150	MEAN
N TIME				
APRIL	3.26	4.18	4.55	4.00
MAY	2.72	3.96	4.29	3.65
MEAN	2.99	4.07	4.42	3.83

STRAW MEAN DM% 80.9

PLOT AREA HARVESTED 0.00355