


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1975

[Full Table of Content](#)



---

### 75/ R/WW/3 - Seed Dressings & Septoria - Winter Wheat

#### Rothamsted Research

Rothamsted Research (1976) 75/ R/WW/3 - *Seed Dressings & Septoria - Winter Wheat* ; Yields Of The Field Experiments 1975, pp 291 - 292 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/WW/3

WINTER WHEAT

SEED DRESSINGS AND SEPTORIA

Object: To study the importance and efficiency of organo-mercury and non-mercury seed dressings against certain pathogens, particularly Septoria seedling blight. Also to study the importance of seedling infections in the epidemiology of the disease - Pastures.

Sponsor: G.L. Bateman.

Design: 2 blocks of 15 plots.

Whole plot dimensions: 2.41 x 11.6.

Treatments: Seed dressings, to seed infected with Septoria (per kg seed):

	SEEDRESS
None (3 plots per block)	0
Phenyl mercury acetate at 1 mg Hg) (2 plots	PMA 1
Phenyl mercury acetate at 5 mg Hg) each per	PMA 5
Phenyl mercury acetate at 25 mg Hg) block)	PMA 25
Carboxin at 4.5 g (a.i.)	CARBOXIN
Guazatine at 0.8 g (a.i.)	GUAZATIN
Guazatine at 0.6 g + maneb at 0.6 g (a.i.)	GUAZ/MAN
Maneb at 3.2 g (a.i.)	MANEB
Quinacetol sulphate at 0.27 g + maneb at 0.45 g (a.i.)	QUIN/MAN
Thiabendazole at 2.4 g (a.i.)	THIABEND

Basal applications: Manures: 'Nitro-Chalk' at 380 kg.

Seed: Chalk, sown at 200 kg.

Cultivations, etc.: - Chisel ploughed twice: 9 Dec, 1974. Rotary harrowed: 19 Dec. Seed sown: 20 Dec. N applied: 21 Apr, 1975. Combine harvested: 18 Aug. Previous crops: Beans 1973, potatoes 1974.

NOTE: Counts were made of seedling emergence. Assessments were made of seedlings infected by Septoria, and on one occasion of leaf infection by Septoria, mildew and eyespot.

75/R./WW/3

GRAIN TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

SEEDRESS	
0	7.27
PMA 1	7.40
PMA 5	7.25
PMA 25	7.08
CARBOXIN	7.16
GUAZATIN	7.01
GUAZ/MAN	7.11
MANEB	7.08
QUIN/MAN	7.30
THIABEND	7.36
MEAN	7.22

\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*

TABLE	SEEDRESS
SED	0.175 (1)
	0.222 (2)
	0.192 (3)
	0.235 (4)
	0.272 (5)

- (1) 0 V ANY OF PMA
- (2) 0 V ANY OF REMAINDER (EXCLUDING 0 AND PMA)
- (3) PMA
- (4) PMA V ANY OF REMAINDER
- (5) ANY OF REMAINDER

\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	19	0.272	3.8

GRAIN MEAN DM% 85.0

PLOT AREA HARVESTED 0.00126