

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1975

[Full Table of Content](#)



---

## 75/ S/B/1 - Varieties, N & Fungicides - Barley

### Rothamsted Research

Rothamsted Research (1976) *75/ S/B/1 - Varieties, N & Fungicides - Barley* ; Yields Of The Field Experiments 1975, pp 331 - 335 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/S/B/1

SPRING BARLEY

VARIETIES, N AND FUNGICIDES

Object: To study the effects of rates, times and forms of nitrogen and of fungicides on the incidence of foliar diseases and yields of two varieties of barley - Saxmundham, Grove Plot.

Sponsors: F.V. Widdowson, J.F. Jenkyn, A. Penny.

Design: Single replicate of 32 plots split into 2.

Whole plot dimensions: 2.43 x 12.2.

Treatments: All combinations of:-

Whole plots: 1. Varieties:

	VARIETY
Julia	JULIA
Wing	WING
2. Solid nitrogen fertiliser (kg N):	S N
50	50
100	100
3. Time of applying solid nitrogen fertiliser:	S N TIME
Seedbed on 28 Apr	SEEDBED
Top dressed on 29 May	TOPDRESS
4. Liquid nitrogen fertiliser (kg N):	L N
None	0
50 (half on 26 June, half on 17 July)	50
5. Mildew control:	MILD CON
None	NONE
Seed dressed ethirimol crop sprayed tridemorph at 0.53 kg in 280 l on 26 June	ETH/TRID

75/S/B/1

Sub plots: 6. Rust control:

RUST CON

None

NONE

Crop sprayed benodanil at 1.12 kg  
in 280 l on 26 June, 17 July

BENODANI

NOTE: 'Nitro-Chalk' was used as solid fertiliser 'Agsol N26' as  
liquid fertiliser in 75 litres.

Basal applications: Manures: (0:20:20) at 310 kg. Weedkiller: Dichlorprop  
plus MCPA ('Mephetol Plus' at 5.6 l in 340 l),

Seed: Sown at 190 kg.

Cultivations, etc.: - Ploughed: 15 Oct, 1974. Seed sown: 28 Apr, 1975.  
Weedkiller applied: 11 June. Combine harvested: 20 Aug. Previous  
crops: Barley 1973 and 1974.

NOTE: Mildew (*Erysiphe graminis*), brown rust (*Puccinia hordei*) and  
yellow rust (*P. striiformis*) were assessed on 29 July.

75/S/B/1

BARLEY

GRAIN TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

S N	50	100	MEAN
VARIETY			
JULIA	3.33	3.66	3.49
WING	3.60	3.86	3.73
MEAN	3.46	3.76	3.61
S N TIME	SEEDBED	TOPDRESS	MEAN
VARIETY			
JULIA	3.70	3.28	3.49
WING	4.09	3.37	3.73
MEAN	3.89	3.33	3.61
S N TIME	SEEDBED	TOPDRESS	MEAN
S N			
50	3.82	3.11	3.46
100	3.97	3.54	3.76
MEAN	3.89	3.33	3.61
L N	0	50	MEAN
VARIETY			
JULIA	3.38	3.60	3.49
WING	3.61	3.84	3.73
MEAN	3.50	3.72	3.61
L N	0	50	MEAN
S N			
50	3.32	3.61	3.46
100	3.63	3.84	3.76
MEAN	3.50	3.72	3.61
L N	0	50	MEAN
S N TIME	SEEDBED	TOPDRESS	MEAN
SEEDBED	3.35	3.94	3.89
TOPDRESS	3.15	3.50	3.33
MEAN	3.50	3.72	3.61
MILD CON	NONE	ETH/TRID	MEAN
VARIETY			
JULIA	3.31	3.67	3.49
WING	3.65	3.81	3.73
MEAN	3.48	3.74	3.61
MILD CON	NONE	ETH/TRID	MEAN
S N			
50	3.31	3.61	3.46
100	3.65	3.86	3.76
MEAN	3.48	3.74	3.61

75/S/B/1

BARLEY

GRAIN TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

MILD CON	NONE	ETH/TRID	MEAN
S N TIME			
SEEDBED	3.81	3.98	3.89
TOPDRESS	3.15	3.50	3.33
MEAN	3.48	3.74	3.61
MILD CON	NONE	ETH/TRID	MEAN
L N			
0	3.39	3.61	3.50
50	3.57	3.87	3.72
MEAN	3.48	3.74	3.61
RUST CON	NONE	BENODANI	MEAN
VARIETY			
JULIA	3.37	3.61	3.49
WING	3.69	3.77	3.73
MEAN	3.53	3.69	3.61
RUST CON	NONE	BENODANI	MEAN
S N			
50	3.34	3.59	3.46
100	3.72	3.79	3.76
MEAN	3.53	3.69	3.61
RUST CON	NONE	BENODANI	MEAN
S N TIME			
SEEDBED	3.73	4.06	3.89
TOPDRESS	3.33	3.32	3.33
MEAN	3.53	3.69	3.61
RUST CON	NONE	BENODANI	MEAN
L N			
0	3.47	3.53	3.50
50	3.59	3.85	3.72
MEAN	3.53	3.69	3.61
RUST CON	NONE	BENODANI	MEAN
MILD CON			
NONE	3.46	3.51	3.48
ETH/TRID	3.60	3.87	3.74
MEAN	3.53	3.69	3.61



75/S/B/1

GRAIN TONNES/HECTARE

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	VARIETY	S N	S N TIME	L N
SED	0.102	0.102	0.102	0.102
TABLE	MILD CON	RUST CON	VARIETY S N	VARIETY S N TIME
SED	0.102	0.122	0.144	0.144
TABLE	S N S N TIME	VARIETY L N	S N L N	S N TIME L N
SED	0.144	0.144	0.144	0.144
TABLE	VARIETY MILD CON	S N MILD CON	S N TIME MILD CON	L N MILD CON
SED	0.144	0.144	0.144	0.144
TABLE	VARIETY RUST CON	S N RUST CON	S N TIME RUST CON	L N RUST CON
SED	0.159	0.159	0.159	0.159
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:				
VARIETY	0.172			
S N		0.172		
S N TIME			0.172	
L N				0.172
TABLE	MILD CON RUST CON			
REP	16			
SED	0.159			
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:				
MILD CON	0.172			

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
WP	6	0.289	8.0
WP.SP	16	0.487	13.5

GRAIN MEAN DM% 82.2

SUB PLOT AREA HARVESTED 0.00073