
Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1975



Full Table of Content

75/ R&w/P/2 - Seed Stocks & Seed Treatment - Potatoes

Rothamsted Research

Rothamsted Research (1976) 75/ R&w/P/2 - Seed Stocks & Seed Treatment - Potatoes; Yields Of The Field Experiments 1975, pp 355 - 360 - DOI: https://doi.org/10.23637/ERADOC-1-141

75/R/P/2 and 75/W/P/2

POTATOES

SEED STOCKS AND SEED TREATMENT

Object: To study the effects of treating tubers with systemic fungicides or a fumigant on tuber-borne diseases and yield of potatoes - Rothamsted (R) Long Hoos I/II and Wcburn (W) Far Field I.

Sponsors: G.A. Hide, M.J. Adams, F. Bell.

Design: 4 randomised blocks of 6 plots split into 4 (plus one extra block for sampling).

Whole plot dimensions: 5.69 x 9.53.

Treatments: All combinations of:-Whole plots: 1. Varieties:

VARIETY

King Edward, Long Hoos I/II (R) only Maris Piper, Far Field I (W) only Pentland Crown Pentland Ivory

EDWARD PIPER CROWN IVORY

2. Seed hoalth:

HEALTH

ex FS, once-grown at Rothamsted in 1974 ex VTSC, once-grown at Rothamsted in 1974

FS VTSC

Sub plots: 3. Fungicide to seed tubers:

FUNGCIDE

None
Benomyl
See-butylamine
Thiabendazole

NONE BENOMYL SEEBUTYL THIABEND

Basal applications:-

Long Hoos I/II (R): Manures: (13:13:20) at 1510 kg. Weedkiller: Linuron at 1.1 kg plus paraquat at 0.42 kg ion in 450 1. Insecticide: Demeton-s-methyl at 0.25 kg in 450 1. Fungicide with insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 450 1.

75/R/P/2 and 75/W/P/2

Far Field I (W): Manures: (13:13:20) at 1880 kg. Weedkiller: Linuron at 1.2 kg plus paraquat at 0.28 kg ion in 280 l. Insecticide: Demeton-s-methyl at 0.25 kg in 280 l. Fungicide: Mancozeb at 1.3 kg in 390 l.

Cultivations, etc.:-

Long Hocs I/II (R): Ploughed: 7 Jan, 1975. Spring-time cultivated: 26 Apr. NPK applied: 1 May. Spring-time cultivated, spike rotary cultivated, potatoes planted: 5 May. Grubbed: 14 May. Rotary ridged: 22 May. Weedkiller applied: 30 May. Insecticide applied: 27 June. Fungicide with insecticide applied: 28 July. Haulm mechanically destroyed: 26 Sept. Sprayed with undiluted BOV at 170 1: 29 Sept. Lifted: 14 Oct. Previous crops: Barley 1973, beans 1974.

Far Field I (W): Ploughed: 9-10 Jan, 1975. NPK applied: 18 Apr. Deep-tine cultivated: 21 Apr. Spring-tine cultivated: 3 May. Rotary harrowed, potatoes planted: 8 May. Weedkiller applied: 30 May. Insecticide applied: 26 June. Fungicide applied: 16 July. Haulm mechanically destroyed: 29 Sept. Sprayed with undiluted BOV at 160 1: 2 Oct. Lifted: 9 Oct. Previous crops: Fallow 1973, beans 1974.

NOTES: (1) Counts of plant and stem numbers were made before burning off.
(2) Crop samples were taken in July and October for tuber weight, size and estimates of fungal infections.

(3) At harvest tubers were graded into 6 sizes and assessments made of Oospora, Rhizoctonia, Helminthosporium and Phoma infection.

75/R/P/2 LONG HOOS I/II (R) TOTAL TUBERS TONNES/HECTARE

*** TABLE OF MEANS ***

HEALTH FS VTSC MEAN VARIETY EDWARD 19.9 17.6 18.7 CROWN 22.6 20.7 21.6 IVORY 17.1 20.8 19.0 MEAN 19.8 19.7 19.8	
EDWARD 19.9 17.6 18.7 CROWN 22.6 20.7 21.6 IVORY 17.1 20.8 19.0	
CROWN 22.6 20.7 21.6 IVORY 17.1 20.8 19.0	
IVORY 17.1 20.8 19.0	
2.4	
MEAN 19.8 19.7 19.8	
FUNGCIDE NONE BENOMYL SECBUTYL THIABEND VARIETY	MEAN
EDWARD 18.3 18.9 20.1 17.7	18.7
CROWN 21.4 21.8 22.3 21.0	21.6
IYORY 18.4 20.2 18.3 18.9	19.0
200 200	2000
MEAN 19.4 20.3 20.3 19.2	19.8
FUNGCIDE NONE BENOMYL SECBUTYL THIABEND	MEAN
HEALTH	
FS 19.9 20.2 20.0 19.3	19.8
VTSC 18.9 20.4 20.5 19.1	19.7
MEAN 19.4 20.3 20.3 19.2	19.8
FUNGCIDE NONE BENOMYL SECBUTYL	THIABEND
VARIETY HEALTH	198
EDWARD FS 20.1 20.4 20.6	18.6
VTSC 16.6 17.3 19.6	16.8
CROW N FS 22.6 23.0 22.2	22.3
VTSC 20.2 20.6 22.4	19.6
IVORY FS 16.9 17.1 17.3	17.0
VTSC 19.9 23.3 19.4	20.8

***** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	VAI	RIETY	HEALTH	FUNGCIDE	VAR LETY HEALTH
SED		2.58	2.10	0.57	3.65
TABLE	-	RIETY	HEALTH FUNGCIDE	VARIETY HEALTH FUNGCIDE	
SED EXCEPT WHEN C	OMPARING	2.72 MEANS 0.99	2.22 WITH SAME LE	3.84 VEL(S) OF:	
HEALTH VARIETY .HEA	ALTH	0.55	0.81	1.41	

75/R/P/2 LONG HOOS I/II (R)

TOTAL TUBERS TONNES/HECTARE

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM	DF	SE	CV %	
BLOCK.WP	15	5.16	26.1	
BLOCK . WP .S P	54	1.99	10.1	

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

*** TABLE OF MEANS ***

HEALTH	FS	VTSC	MEAN		
VARIETY					
EDWARD	35.9	22.1	29.0		
CROWN	74.9	69.9	72.4		
IVORY	80.9	77.2	79.0		
MEAN	63.9	56.4	60.1		
FUNGCIDE	NONE	BENOM YL	SECBUTYL	THIABEND	MEAN
VARIETY					
EDWARD	30.1	27.2	29.0	29.7	29.0
CROWN	71.0	71.8	77.9	68.8	72.4
IVORY	78.0	79.0	79.0	80.2	79.0
MEAN	59.7	59.3	62.0	59.6	60.1
FUNGCIDE HEALTH	NONE	BENOMYL	SECBUTYL	THIABEND	MEAN
FS	64.5	64.2	63.6	63.3	63.9
VTSC	54.9	54.5	60.3	55.8	56.4
1130	04.9	04.0	00.5	33.0	30.4
MEA N	59.7	59.3	62.0	59.6	60.1
	FUNGCIDE	E NO	NE BENOM	YL SECBUTYL	THIABEND
VARIETY	HEALTH	ł			
ED WARD	FS	37	•1 36	.0 33.3	37.3
	VTSC	23	.1 18	.4 24.7	22.1
CROWN	FS	75	.4 75	.3 77.6	71.3
	VTSC	66		.3 78.1	66.3
IVORY	FS	81	.2 81	.2 80.0	81.3
	VTSC			.8 78.1	79.1

PLOT AREA HARVESTED 0.00135

75/W/P/2 FAR FIELD I (W)

TOTAL TUBERS TONNES / HECTARE

*** TABLE OF MEANS ***

HEALTH	FS	VTSC	MEAN		
VARIETY					
PIPER	19.4	18.3	18.9		
CROWN	32.9	33.5	33.2		
IVORY	22.4	23.2	22.8		
MEAN	24.9	25.0	25.0		
FUNGCIDE VARIETY	NONE	BENOMYL	SECHUTYL	THI ABEND	MEAN
PIPER	17.5	19.1	18.4	20.6	18.9
CROWN	34.2	31.5	32.7	34.3	33.2
IVORY	20.1	22.2	25.5	23.4	22.8
MEAN	23.9	24.2	25.5	26.1	25.0
FUNGCIDE HEALTH	NONE	BENOMYL	SECBUTYL	THIABEND	MEAN
FS	22.7	25.7	25.7	25.5	24.9
VISC	25.1	22.8	25.3	26.7	25.0
412C	20.1	22.0	20.0	20.1	20.0
MEAN	23.9	24.2	25.5	26.1	25.0
	FUNGCIDE	NO NO	NE BENOM	YL SECBUTYL	THIABEND
VARIETY	HEALTH	I			
PIPER	FS	17	•1 19	.9 19.5	21.3
	VTSC	17	.8 18	.3 17.3	20.0
CROWN	FS	31	•5 32	.8 33.0	34.2
	VTSC	36	.9 30	.3 32.4	34.3
IVORY	FS	19	.6 24	.5 24.6	20.9

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE	VAI	RIETY	HEA	LTH	FUNG	CIDE	VARIETY HEALTH
SED		1.25		.02		0.99	1.76
TABLE		RIETY	HEA FUNGO	LTH		IETY ALTH CIDE	
SED EXCEPT WHEN VARIETY	COMPARING	1.94 MEANS 1.71	and the second second second second	.58 ME LE	The second second	2.74 OF:	
HEALTH VARIETY .H	EALTH		1	1.40		2.42	

75/4/P/2 FAR FIELD I (W)

TOTAL TUBERS TONNES/HECTARE

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLCCK.WP	15	2.49	10.0
BLCCK.WP.SP	54	3.43	13.7

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

*** TABLES OF MEANS ***

HEALTH	FS	VTSC	MEA	N		
VARIETY		22.5	0.5	_		
PIPER	28.1	22.8	25.			
CROWN	78.2	74.0	76.			
IVORY	74.9	70.1	72.	5		
MEAN	60.4	55.6	58.	0		
FUNGCIDE VARIETY	NONE	BENOMYL	SECBUTY	L TH	IABEND	MEAN
PIPER	24.1	26.8	23.	6	27.3	25.5
CROWN	74.5	74.9	78.	5	76.6	76.1
IVORY	69.2	70.2	75.	5	75.1	72.5
MEAN	55.9	57.3	59.	2	59.7	58.0
FUNGCIDE	NONE	BENOMY L	SECBUTY	L TH	IABEND	MEAN
HEALTH	F0 0	CO 4		-	60.0	CO 4
FS	59.0	60.4	61.		60.9	60.4
VTSC	52.9	54.2	57.	1	58.4	55.6
MEAN	55.9	57.3	59.	2	59.7	58.0
	FUNGCID	E NO	NE BEN	OMYL	SECBUTYL	THIABEND
VARIETY	HEALT	H				
PIPER	F	S 24	•3	29.6	28.2	30.3
	VTS	C 24	.0	24.0	18.9	24.3
CROWN	F	S 79	0.1	77.1	77.6	78.8
	VTS	C 69	9.9	72.7	79.3	74.4
I VORY	F	S 73	5.5	74.4	78.1	73.6
	VTS	C 64	.9	65.9	73.0	76.5

PLOT AREA HARVES TED 0.00135