


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1975

[Full Table of Content](#)



### 75/ R/ON/1 - Stem-eelworm Control - Onions

#### Rothamsted Research

Rothamsted Research (1976) 75/ R/ON/1 - *Stem-eelworm Control - Onions* ; Yields Of The Field Experiments 1975, pp 375 - 376 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/ON/1

WINTER AND SPRING ONIONS

STEM-EELWORM CONTROL

Object: To study the effects of rates and times of applying aldicarb on control of stem-eelworm (*Ditylenchus dipsaci*) and on the yield of winter onions. Spring-sown onions treated with aldicarb are also included - Great Field II.

Sponsor: A.G. Whitehead.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 1.22 x 5.18.

Treatments: Aldicarb (kg):-

ALDICARB

To winter onions:

None (2 plots/block)	0
1 in seedbed, 1 in spring	1+1
2 in seedbed, none in spring	2
2 in seedbed, 2 in spring	2+2
4 in seedbed, none in spring	4
4 in seedbed, 4 in spring	4+4
8 in seedbed, none in spring	8

To spring onions:

4 kg in seedbed (2 plots/block)

NOTE: Yields were not taken from the spring-sown onions.

Standard applications: Manures: Autumn-sown plots: (13:13:20) at 1040 kg, 'Nitro-Chalk' at 450 kg. Spring-sown plots: (13:13:20) at 1880 kg. Weedkiller: Propachlor ('Ramrod' at 6.7 kg in 450 l).

Seed: Winter: Imai Early Yellow, dressed with dieldrin, sown at 6.7 kg. Spring: Robusta, dressed with dieldrin, sown at 6.7 kg.

Cultivations, etc.:- Ploughed: 22 Aug, 1974. Autumn-sown plots: rotary cultivated twice, NPK and aldicarb applied, seed sown: 22 Aug. Weedkiller applied: 27 Aug. N applied: 14 Feb, 1975. Aldicarb applied: 21 Apr. Lifted: 4 July. Spring-sown plots: Weedkiller applied: 27 Aug, 1974. Rotary cultivated, NPK and aldicarb applied and seed sown: 21 Apr, 1975. Previous crops: Onions 1973, barley 1974.

NOTES: (1) Soil samples were taken from winter-sown plots on 23 July for counts of *Ditylenchus dipsaci*.  
(2) Onions were stored until 3 October to determine loss of weight and percentage of bulbs rotting after harvest.

75/R/ON/1

SALEABLE ONIONS TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

ALDICARB	0	1+1	2	2+2	4	4+4	8	MEAN
	13.2	18.6	18.8	19.3	18.5	19.2	19.3	17.5

\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*

TABLE ALDICARB

-----	-----
SED	2.74(1)
	3.17(2)
(1) 0 V ANY OF REMAINDER	
(2) ANY OF REMAINDER	

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	15	3.88	22.1

PLOT AREA HARVESTED 0.00063