


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1975

[Full Table of Content](#)



## 75/ R/RN/8 - Cultivation/WEEDKILLER - Barley

### Rothamsted Research

Rothamsted Research (1976) 75/ R/RN/8 - Cultivation/WEEDKILLER - Barley ; Yields Of The Field Experiments 1975, pp 97 - 99 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/RN/8

CULTIVATION/WEEDKILLER

Object: To study the long term effects of weedkillers and different methods of primary cultivation on a rotation of crops - Great Harpenden I.

Sponsors: R. Moffitt, G.V. Dyke, J.A. Currie.

The 15th year, barley.

For previous years see 'Details' 1967, 68/E/6(t), 69/R/RN/8(t), 70/R/RN/8, 71/R/RN/8(t) and 72-74/R/RN/8.

Design: 2 randomised blocks of 12 plots split into 2.

Whole plot dimensions: 12.8 x 15.2.

Treatments: All combinations of:-

Whole plots: 1. Primary cultivations annually:

Ploughed, 19 Mar, 1975  
Rotary cultivated, 20 Mar  
Deep-tine cultivated twice, 17, 18 Mar

CULTIVAT

PLOUGH  
ROTAVATE  
DEEPTINE

2. Weed control to potatoes 1974:

WEEDCNTR(74)

Mechanical  
Linuron plus paraquat  
Linuron plus paraquat. Grubbed and rotary ridged

MECHANCL  
LIN/PAR  
LIN/PARR

Sub plots: 3. Hormone weedkiller to barley 1975:

WEEDKILL(75)

None  
Dicamba plus mecoprop and MCPA ('Tetralix Plus'  
at 7.0 l in 220 l) 6 June, 1975

NONE

DI/ME/MC

4. Paraquat weedkiller to wheat stubble  
autumn 1973:

WEEDKILL(73)

None  
Paraquat

NONE

PARAQUAT

plus three extra treatments:-

EXTRA

75/R/RN/8

Spike rotary cultivated, 23 Apr. Given linuron plus paraquat to potatoes 1974, with sub plot treatments 3 and 4 above.

SPIKE

Shallow ploughed, 28 Feb. Given linuron plus paraquat to potatoes 1974, whole plot given paraquat as 4 above, with sub plot treatment 3 above

(SH)PLCH

Standard cultivations as considered best for each crop. Deep-tine cultivated twice, 19 Mar for barley. Given linuron plus paraquat and grubbed and rotary ridged to potatoes 1974, with sub plot treatments 3 and 4 above.

STANDARD

Basal applications: Manures (20:14:14) at 380 kg combine drilled.

Seed: Julia, dressed with ethirimol, sown at 160 kg.

Cultivations, etc.:- Spring-tine cultivated: 24 Dec, 1974. Seed sown: 23 Apr, 1975. Combine harvested: 21 Aug.

75/R/RN/3

BARLEY

GRAIN TONNES/HECTARE

\*\*\* TABLES OF MEANS \*\*\*

OMITTING EXTRA PLOTS

CULTIVTN	PLOUGH	ROTAVATE	DEEPTINE	MEAN
WEEDCNTL(74)				
MECHANCL	3.70	3.33	3.68	3.57
LIN/PAR	3.79	3.55	3.72	3.69
LIN/PARR	3.83	3.43	4.09	3.78
WEEDKLLR(75)				
NONE	3.89	3.74	3.54	3.72
DI/ME/MC	3.66	3.14	4.12	3.64
WEEDKLLR(73)				
NONE	3.94	3.56	3.73	3.74
PARAQUAT	3.61	3.32	3.94	3.62
MEAN	3.77	3.44	3.83	3.68

EXTRA PLOTS ONLY

EXTRA	SPIKE	(SH)PLGH	STANDARD
WEEDKLLR(75)			
NONE	3.53	4.06	3.89
DI/ME/MC	3.80	4.15	4.51
WEEDKLLR(73)			
NONE	4.04		4.00
PARAQUAT	3.29	4.11	4.39
MEAN	3.66	4.11	4.20

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	CULTIVTN	WEEDCNTL(74)	WEEDKLLR(73)	WEEDKLLR(75)
SED	0.189	0.189	0.143	0.143

TABLE	CULTIVTN	CULTIVTN	CULTIVTN
	WEEDCNTL(74)	WEEDKLLR(73)	WEEDKLLR(75)
SED	0.328	0.258	0.258
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:			
CULTIVTN		0.248	0.248

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	8	0.328	8.9
BLOCK.WP.SP	8	0.429	11.7

GRAND MEAN 3.76

GRAIN MEAN DM% 81.6

SUB PLOT AREA HARVESTED 0.00434