Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1976



Full Table of Content

76/W/RN/16 Effects of Deep P K - Wheat, Sugar Beet, Barley, Potatoes

Rothamsted Research

Rothamsted Research (1977) 76/W/RN/16 Effects of Deep P K - Wheat, Sugar Beet, Barley, Potatoes ; Yields Of The Field Experiments 1976, pp 112 - 116 - **DOI:**

https://doi.org/10.23637/ERADOC-1-15

EFFECTS OF DEEP PK

Object: To study the residual effects of subsoiling, and of incorporating a large dressing of PK in either the subsoil or topsoil, on yields of a rotation of crops - Woburn Butt Furlong.

Sponsor: J. McEwen.

The third year, winter wheat, sugar beet, spring barley, potatoes.

For previous year see 74-75/W/RN/16.

Design: 4 series of 3 randomised blocks of 4 plots.

Whole plot dimensions: 4.27 x 2.59.

Treatments: Extra PK and subsoil treatment (applied autumn 1973):

PK SUB Extra PK Subsoil (25-50 cm) treatment

- None None Subsoiled

PKTOP - To topsoil (0-25 cm) None

- PKSUB To subsoil

- NOTES: (1) The rates of P and K were 1930 kg P205, as superphosphate and 460 kg K20 as muriate of potash. These quantities, applied to subsoil, were chosen to equalize available P and K in top and subsoil.
 - (2) Subsoiling was done by spade, after removing the topsoil which was then replaced. PK to subsoil was worked in by forking.
 - (3) PK to topsoil was applied half before ploughing in autumn half soon after on the plough furrow.

Standard applications:

Series III: Winter wheat: Manures: (0:20:20) at 290 kg, combine drilled.

N at 75 kg as 'Nitro-Chalk'. Weedkiller: Ioxynil at 0.6 kg plus
mecoprop at 1.8 kg in 340 l. Insecticide: Pirimicarb at 0.14 kg in 280 l.

Series IV: Sugar beet: Manures: Magnesian limestone at 5 tonnes.
(0:14:28) at 750 kg. N at 140 kg as 'Nitro-Chalk'. Insecticide:
Pirimicarb at 0.14 kg in 280 l.

Series I: Barley: Manures: (20:14:14) at 380 kg, combined drilled. Weedkiller: Ioxynil at 0.6 kg plus mecoprop at 1.8 kg in 340 l. Insecticide: Pirimicarb at 0.14 kg in 280 l.

Series II: Potatoes: Manures: (13:13:20) at 1860 kg. Weedkiller: Linuron at 0.9 kg in 430 l. Insecticide: Pirimicarb at 0.14 kg in 280 l.

Seed: Winter wheat: Cappelle, sown at 210 kg.
Sugar beet: Klein E, sown at 5.6 kg.
Barley: Julia, dressed with ethirimol, sown at 160 kg.
Potatoes: Pentland Crown.

Cultivations, etc .:-

Series III: Winter wheat: Spring-tine cultivated with crumbler, seed sown: 23 Oct, 1975. N applied: 13 Apr, 1976. Weedkiller applied: 30 Apr. Insecticide applied: 25 June. Hand harvested: 19 July. Series IV: Sugar beet: Magnesian limestone applied: 5 Sept, 1975. Ploughed: 28 Oct. N and PK applied: 30 Mar, 1976. Spring-tine cultivated with crumbler twice, seed sown: 5 Apr. Singled by hand 22 May. Insecticide applied: 25 June. Hand lifted: 17 Nov. Series I: Barley: Ploughed: 24 Nov, 1975. Spring-tine cultivated with crumbler: 8 Mar, 1976. Seed sown: 9 Mar. Weedkiller applied: 30 Apr. Insecticide applied: 25 June. Hand harvested: 20 July. Series II: Potatoes: Ploughed: 28 Oct, 1975. NPK applied: 23 Mar, 1976. Rotary harrowed, potatoes planted: 31 Mar. Weedkiller applied: 30 Apr. Insecticide applied: 25 June. Hand lifted: 19 Oct.

NOTE: Samples of wheat and barley grain, potato tubers and sugar beet roots and tops were taken for analysis of N,P,K,Na,Ca and Mg.

WINTER WHEAT

GRAIN TONNES/HECTARE

**** TABLES OF MEANS ****

PK SUB -- - SUB PKTOP - - PKSUB MEAN 3.48 4.06 3.18 4.17 3.72

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE PK SUB
SED 0.281

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV%

BLOCK.WP 6 0.344 9.2

GRAIN MEAN DM% 89.2

STRAW TONNES/HECTARE

**** TABLES OF MEANS ****

PK SUB -- - SUB PKTOP - - PKSUB MEAN 4.90 5.64 4.33 6.34 5.30

STRAW MEAN DM% 90.3

PLOT AREA HARVESTED 0.00033

SUGAR BEET

ROOTS WASHED TONNES/HECTARE

***** TABLES OF MEANS ****

PK SUB -- - SUB PKTOP - - PKSUB MEAN 35.3 36.9 33.0 39.3 36.1

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE PK SUB
SED 1.62

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

 STRATUM
 DF
 SE
 CV%

 BLOCK.WP
 6
 1.98
 5.5

SUGAR BEET

SUGAR PERCENTAGE TONNES/HECTARE

**** TABLES OF MEANS ****

PK SUB -- - SUB PKTOP - - PKSUB MEAN 14.9 15.0 14.8 14.9 14.9

TOTAL SUGAR TONNES/HECTARE

**** TABLES OF MEANS ****

PK SUB -- - SUB PKTOP - - PKSUB MEAN 5.25 5.54 4.88 5.86 5.38

***** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE PK SUB
SED 0.241

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

 STRATUM
 DF
 SE
 CV%

 BLOCK.WP
 6
 0.295
 5.5

TOPS TONNES/HECTARE

***** TABLES OF MEANS ****

PK SUB -- - SUB PKTOP - - PKSUB MEAN 28.3 24.9 24.7 25.6 25.9

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE PK SUB
SED 1.32

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

 STRATUM
 DF
 SE
 CV%

 BLOCK.WP
 6
 1.62
 6.3

PLOT AREA HARVESTED 0.00049

BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS ****

PK SUB -- SUB PKTOP - PKSUB MEAN 3.43 4.46 3.77 4.51 4.04

**** STANDARD ERRORS OF DIFFERENCES OF MEANS ****

TABLE PK SUB
SED 0.261

**** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ****

STRATUM DF SE CV%

BLOCK.WP 6 0.319 7.9

GRAIN MEAN DM% 89.2

STRAW TONNES/HECTARE

**** TABLES OF MEANS ****

PK SUB -- - SUB PKTOP - - PKSUB MEAN 3.16 4.14 3.41 4.44 3.79

STRAW MEAN DM% 92.2

PLOT AREA HARVESTED 0.00033

POTATOES

TOTAL TUBERS TONNES/HECTARE

**** TABLES OF MEANS ****

PK SUB -- - SUB PKTOP - - PKSUB MEAN 41.3 39.7 43.0 47.1 42.8

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE PK SUB
SED 2.34

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

 STRATUM
 DF
 SE
 CV%

 BLOCK.WP
 6
 2.87
 6.7

PLOT AREA HARVESTED 0.00043