


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1976

[Full Table of Content](#)



### 76/R/CS/41 Cultivations and Soil Invertebrates - Old Grass, New Grass

#### Rothamsted Research

Rothamsted Research (1977) *76/R/CS/41 Cultivations and Soil Invertebrates - Old Grass, New Grass* ; Yields Of The Field Experiments 1976, pp 165 - 167 - DOI:  
<https://doi.org/10.23637/ERADOC-1-15>

76/R/CS/41

CULTIVATIONS AND SOIL INVERTEBRATES

Object: To study the effects of cultivations on yields of grass and on populations of soil animals - Road Piece.

Sponsor: C.A. Edwards.

The eighth year, old grass, new grass.

For previous years see 69/R/CS/41(t), 70/R/CS/41(t) and 71-75/R/CS/41.

Design: 4 blocks of 8 plots, randomisation restricted.

Whole plot dimensions: 6.40 x 7.32.

Treatments: Cultivations and reseeded:

CULTIVTN

- 0 No treatments to old grass (two plots per block)  
Grass ploughed up:-
- SF In spring 1969, reseeded after fewest cultivations needed to produce a seedbed
- SM In spring 1969, reseeded after many seedbed cultivations
- AM In autumn 1969, reseeded spring 1970 after many seedbed cultivations
- SFR Every spring since 1969, reseeded each year after fewest cultivations needed to produce a seedbed
- SMR Every spring since 1969, reseeded each year after many seedbed cultivations
- AMR Every autumn since 1969, reseeded every following spring after many seedbed cultivations

Seeds mixture for 1976: S215 meadow fescue at 9.4 kg; Contessa meadow fescue at 6.0 kg; Pecora certified timothy at 6.0 kg; N.Z. Huia white-clover at 2.2 kg; Wild white clover at 0.4 kg. Sown at 24 kg.

Basal applications: Manures: (0:14:28) at 500 kg in winter. (25:0:16) at 440 kg in spring and at 220 kg after the first cut.

Cultivations, etc.: - Basal PK applied: 21 Nov, 1975. AMR plots ploughed: 11 Dec. SMR and SFR plots ploughed: 9 Feb, 1976. Basal NK applied, disced all plots to be reseeded: 26 Feb. Disced AMR and SMR plots four times, SFR plots twice: 13 Apr. Power harrowed and seed sown: 21 Apr. Cut (excluding AMR, SMR, SFR plots): 20 May, 29 June, 2 Nov. NK applied to all cut plots: 21 May.

NOTE: Soil cores were taken to assess total soil fauna in January, March and September. Quadrats were sampled on each plot for earthworms in December 1975, March, 1976 and September.

76/R/CS/41

1ST CUT (20/5/76) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CULTIVTN	0	SF	SM	AM	MEAN
	3.74	3.40	3.00	3.71	3.52

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	CULTIVTN
-----	-----
SED	0.475 MIN REP 0.412 MAX-MIN

CULTIVTN  
 MAX-MIN O V ANY OF REMAINDER  
 MIN REP ANY OF REMAINDER

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	13	0.672	19.1

1ST CUT MEAN DM% 18.8

2ND CUT (29/6/76) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CULTIVTN	0	SF	SM	AM	MEAN
	0.58	0.81	1.28	1.14	0.88

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	CULTIVTN
-----	-----
SED	0.246 MIN REP 0.213 MAX-MIN

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	13	0.348	39.7

2ND CUT MEAN DM% 35.2

76/R/CS/41

3RD CUT (2/11/76) DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CULTIVTN	0	SF	SM	AM	MEAN
	1.73	1.89	1.58	1.55	1.70

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	CULTIVTN
SED	0.117 MIN REP 0.101 MAX-MIN

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	13	0.165	9.7

3RD CUT MEAN DM% 13.5

TOTAL OF 3 CUTS DRY MATTER TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CULTIVTN	0	SF	SM	AM	MEAN
	6.05	6.11	5.86	6.39	6.09

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	CULTIVTN
SED	0.561 MIN REP 0.486 MAX-MIN

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	13	0.794	13.0

TOTAL OF 3 CUTS MEAN DM% 22.5

PLOT AREA HARVESTED 0.00074