


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1976

[Full Table of Content](#)



76/R/CS/90 Cultivation for Cereals - Wheat

Rothamsted Research

Rothamsted Research (1977) *76/R/CS/90 Cultivation for Cereals - Wheat ; Yields Of The Field Experiments 1976*, pp 170 - 171 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/R/CS/90

CULTIVATIONS FOR CEREALS

Object: To study the engineering aspects - power requirements, rate of work, revenue and costs - of different tillage systems for continuous wheat. Effects on weeds, soil pathogens and yields are also studied - Meadow.

Sponsors: D.E. Patterson (N.I.A.E.), R. Moffitt, R.D. Prew.

The fifth year, winter wheat.

For previous years see 72-75/R/CS/90.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 13.7 x 33.8.

Treatments

TILLAGE Tillage systems:-

- 1 Three passages of the tractor (three-pass system): Ploughed* 20 cm deep (8 inches): finger-tine harrowed: drilled.
- 2 Five-pass system: Tine cultivated* 15 cm deep (6 inches): tine cultivated 15 cm: finger-tine harrowed: drilled.
- 3 Two-pass system: Ploughed* 20 cm deep: finger-tine harrowed and drilled.
- 4 Two-pass system: Ploughed* 10 cm deep: finger-tine harrowed and drilled.
- 5 Three-pass system: Tine cultivated* (N.I.A.E.) 20 cm deep: disc harrowed: finger-tine harrowed and drilled.
- 6 Three-pass system: Tine cultivated* (N.I.A.E.) 10 cm deep: disc harrowed: finger-tine harrowed and drilled.
- 7 Two-pass system (stubble of 1975 crop burnt on 4 Sept): Tine cultivated* (N.I.A.E.) 10 cm deep: finger-tine harrowed and drilled.
- 8 Two-pass system: Sprayed with paraquat (0.56 kg ion in 220 l on 14 Oct): direct drilled.
- 9 Three-pass system: Rotary digger* (N.I.A.E.) cultivated: disc harrowed: finger-tine harrowed and drilled.
- 10 Two-pass system (stubble of 1975 crop burnt on 4 Sept): Rotary digger* (N.I.A.E.) cultivated: finger-tine harrowed and drilled.

NOTE: Rotary digger (N.I.A.E) - depth of working: rotor 10 cm, tines 20 cm.

* Cultivations done on 8 Oct, 1975. All other cultivations and drilling done on 15 Oct. A disc drill was used on all treatments except 8.

Basal applications: Manures: (10:24:24) at 310 kg, combine drilled. 'Nitro-Chalk' at 380 kg. Weedkillers: Dicamba with mecoprop and MCPA ('Banlene Plus' at 5.6 l in 220 l). Paraquat at 0.42 kg ion in 220 l. Difenzoquat at 1.0 kg in 220 l. Insecticide: Pirimicarb at 0.14 kg in 220 l.

76/R/CS/90

Seed: Atou, sown at 170 kg.

Cultivations, etc.:- Basal paraquat applied: 2 Oct, 1975. Seed sown: 15 Oct. N applied: 1 Apr, 1976. 'Banlene Plus' applied: 8 Apr. Difenzoquat applied: 28 Apr. Insecticide applied: 23 June. Combine harvested: 31 July.

NOTES: Observations and determinations were made as follows:-

- (1) Soil: Structure assessments, photographs for aggregate size, moisture determinations.
- (2) Implements: Depth and width of work, forward speed, wheel slip and power requirements.
- (3) Crop: Plant and tiller counts, numbers of ears and grains per ear, disease and pest assessments, aerial photographs.
- (4) On one of the blocks, plots of TILLAGE 6 and 9 were very severely infested with sterile brome grass (*Bromus sterilis*). This infestation is thought to have markedly lessened yields from these plots: the yields presented have not been adjusted.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TILLAGE	1	2	3	4	5	6	7	8	9	10	MEAN
	4.01	4.05	3.93	4.08	3.89	3.70	3.90	2.94	3.84	4.04	3.84

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TILLAGE
-----	-----
SED	0.285

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	18	0.349	9.1

GRAIN MEAN DM% 88.0

PLOT AREA HARVESTED 0.01031