

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1976

[Full Table of Content](#)

76/W/CS/99 Effects of Breaks on Take-all - Barley, S. Beans

Rothamsted Research

Rothamsted Research (1977) *76/W/CS/99 Effects of Breaks on Take-all - Barley, S. Beans* ; Yields Of The Field Experiments 1976, pp 172 - 173 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/W/CS/99

EFFECTS OF BREAKS ON TAKE-ALL

Object: To study the phenomenon of 'take-all' (*Gaeumannomyces graminis*) decline in barley - Woburn Butt Furlong.

Sponsor: D. Hornby.

The fifth year, barley, spring beans.

For previous years see 72/W/CS/99(t) and 73-75/W/CS/99.

Design: 2 blocks of 9 plots.

Whole plot dimensions: 5.33 x 15.2.

Treatments: Previous crops:-

PREVCROP	1968-71	1972	1973	1974	1975
B/B/B/B	B	B	B	B	B (3 plots/block)
F/B/B/B	B	F	B	B	B
F/BE/B/B	B	F	BE	B	B
B/F/BE/B	B	B	F	BE	B
B/B/F/BE	B	B	B	F	BE

B = Barley, BE = Beans, F = Fallow

Standard applications:

Barley: (20:14:14) at 420 kg combine drilled. Weedkiller: Ioxynil at 0.53 kg and mecoprop at 1.6 kg in 280 l.
Spring beans: (0:14:28) at 400 kg.

Seed: Barley: Julia, dressed with ethirimol, sown at 160 kg.
Spring beans: Minden, sown at 220 kg.

Cultivations, etc.:-

All plots: Ploughed: 6 Oct, 1975. Spring-tine cultivated: 1 Mar, 1976.
Barley: Seed sown: 3 Mar, 1976. Weedkiller applied: 29 Apr.
Combine harvested: 27 July.
Spring beans: Spring-tine cultivated: 4 Mar, 1976. Seed sown: 11 Mar.
PK applied by hand: 12 Mar. Hoed by hand three times: 28 Apr, 26 May, 22 June. Combine harvested: 20 July.
Fallow: Spring-tine cultivated: 4 Mar, 1976. Hoed by hand twice: 28 Apr, 26 May.

- NOTES: (1) Soil samples were taken before sowing and after harvest and plant samples in July for incidence of 'take-all' (*Gaeumannomyces graminis*).
(2) Yields were not taken for the spring beans.

76/W/CS/99

BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	B/B/B/B	F/B/B/B	F/BE/B/B	B/F/BE/B	B/B/F/BE	MEAN
	1.61	2.01	1.79	2.37	2.85	1.98

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PREVCROP	
-----	-----	
SED	0.459	MIN REP
	0.375	MAX-MIN

PREVCROP
 MAX-MIN B/B/B/B V ANY OF REMAINDER
 MIN-REP ANY OF REMAINDER

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	8	0.459	23.2

GRAIN MEAN DM% 89.2

PLOT AREA HARVESTED 0.00434