

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1976

[Full Table of Content](#)

### 76/W/CS/174 Sowing Dates and Ccn - W. Wheat, S. Wheat, Barley and Oats

#### Rothamsted Research

Rothamsted Research (1977) *76/W/CS/174 Sowing Dates and Ccn - W. Wheat, S. Wheat, Barley and Oats* ; Yields Of The Field Experiments 1976, pp 219 - 221 - DOI:

<https://doi.org/10.23637/ERADOC-1-15>

76/W/CS/174

SOWING DATES AND CCN

Object: To study the effects of sowing date and a nematicide on the incidence of cereal cyst-nematode (*Heterodera avenae*) (CCN) and the yield of three cereals in a soil known to contain a fungal parasite of the nematode - Woburn, Butt Close.

Sponsor: B.R. Kerry.

The second year, winter and spring wheat, barley and oats.

For previous year see 75/W/M/1.

Design: 4 randomised blocks of 12 whole plots, 6 of them split into 2.

Whole plot dimensions: 2.13 x 6.70.

Treatments: All combinations of:-

Whole plots

1. CROP Crop (cumulative to 1975):

WHEAT	Wheat
BARLEY	Barley
OATS	Oats

2. SOW DATE Sowing date (cumulative to 1975):

AUTUMN	Autumn
SPRING	Spring

3. NEMACIDE Nematicide (cumulative to 1975):

NONE	None
OXAMYL	Oxamyl at 8.8 kg

Sub plots

4. STERILNT Sterilant (1976 only):  
(combinations  
with SOW DATE  
SPRING only)

NONE	None
FORMALIN	Formalin at 3000 l in 109000 l

Standard applications:

Autumn-sown cereals: Manures: (10:24:24) at 260 kg, combine drilled.  
Winter wheat: N at 100 kg, as 'Nitro-Chalk'. Winter barley and oats:  
N at 80 kg, as 'Nitro-Chalk'.

76/W/CS/174

Spring-sown cereals: Manures: (20:14:14) at 380 kg combine drilled.  
Fungicide to barley only: Tridemorph at 0.53 kg in 340 l.  
All cereals: Weedkiller: Ioxynil at 0.6 kg plus mecoprop at 1.8 kg  
in 340 l. Insecticide: Pirimicarb at 0.14 kg in 280 l.

Seed: Wheat: Flinor, sown at 210 kg in autumn and 200 kg in spring.  
Barley: Maris Otter, sown at 190 kg in autumn and 160 kg in spring.  
Oats: Peniarth, sown at 200 kg in autumn and 190 kg in spring.

Cultivations, etc.:-

All Cereals: Heavy spring-tine cultivated: 30 Aug, 1975. Rotary  
cultivated: 1 Sept. Ploughed: 24 Sept. Spring-tine cultivated with  
crumbler: 26 Sept. Weedkiller applied: 30 Apr. 1976. Insecticide  
applied: 29 June. Combine harvested: 28 July.

Autumn-sown Cereals: Oxamyl applied and these plots only rotary cultivated:  
2 Oct, 1975. Seed sown: 15 Oct. N applied: 12 Apr, 1976.

Spring-sown Cereals: Spring-tine cultivated twice: 14 Oct, 15 Oct, 1975.  
Formalin applied: 1 Mar, 1976. Oxamyl applied and these plots only  
rotary cultivated, all plots spring-tine cultivated with crumbler:  
22 Mar. Seed sown: 23 Mar. Fungicide applied to barley only: 9 June.

- NOTES: (1) Soil samples were taken before sowing in autumn and spring and  
after harvest for egg counts of *Heterodera avenae*.  
(2) Counts of white female *H.avenae* were made during the growing  
season and amounts of infection of these by *Entomophthora* spp.  
and *Verticillium chlamyosporium* were estimated.  
(3) Yields from SOW DATE SPRING were very small on all plots and  
unrecordable on some, because of drought. No yields are  
presented for combinations with this level.

76/W/CS/174

SOW DATE AUTUMN ONLY

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CROP NEMACIDE	WHEAT	BARLEY	OATS	MEAN
NONE	1.58	1.09	0.73	1.13
OXAMYL	2.40	2.65	2.19	2.41
MEAN	1.99	1.87	1.46	1.77

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	NEMACIDE	CROP	NEMACIDE CROP
SED	0.163	0.199	0.282

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	10	0.345	19.5
GRAIN MEAN DM%	87.0		

STRAW TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CROP NEMACIDE	WHEAT	BARLEY	OATS	MEAN
NONE	1.68	1.23	0.75	1.22
OXAMYL	2.30	1.94	1.51	1.92
MEAN	1.99	1.59	1.13	1.57

STRAW MEAN DM% 85.9

PLOT AREA HARVESTED 0.00095