

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1976

[Full Table of Content](#)



## 76/S/CS/1 Varieties, N and Ccc - Wheat

### Rothamsted Research

Rothamsted Research (1977) *76/S/CS/1 Varieties, N and Ccc - Wheat* ; Yields Of The Field Experiments 1976, pp 248 - 250 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/S/CS/1

VARIETIES, N AND CCC

Object: To study the effects of nitrogen fertiliser, at a range of rates and times, and chlormequat (CCC) on the yield of two varieties of winter wheat - Saxmundham, Oldershaw's and Garner's plots.

Sponsors: F.V. Widdowson, A.E. Johnston.

The 11th year, winter wheat.

For previous years see 66/C/30(t), 67/C/23(t), 68/C/39, 69-70/S/CS/1, 71/S/CS/1(t), 72/S/CS/1(t) and 73-75/S/CS/1.

Design: A single replicate of 4 x 2 x 2 x 2 in 4 blocks of 4 plots, each split lengthways into 2, plus one additional plot per block. Additionally all the plots are split breadthways into 3.

Whole plot dimensions: 5.49 x 40.2.

Treatments: All combinations of:-

Whole plots (All sown at a seed rate of 170 kg with 13 cm (5 inches) between the rows):

1. PREVCROP      Number of previous continuous wheat crops:

7 WHEAT	7
8 WHEAT	8
9 WHEAT	9
10 WHEAT	10

2. CCC      Chlormequat (kg):

0.0	None
1.7	1.7 in 340 l

Half plots

3. N TIME      Times of applying nitrogen fertiliser:

SINGLE	Single dressing (21 Apr)
DIVIDED	Divided dressing (Half 21 Apr, half 18 May)

4. VARIETY      Varieties:

CAPPELLE	Cappelle
HUNTSMAN	Maris Huntsman

Pairs of sixth plots

5. N RATE      Rates of nitrogen fertiliser in addition to 62 kg N in autumn (14 Oct) (kg N):

50	50
100	100
150	150

76/S/CS/1

EXTRA Together with one extra plot per block which had 6 previous wheat crops and was sown with Cappelle at a seed rate of 180 kg with 20 cm (8 inches) between the rows and tested all combinations of:-

Half plots

1. AUT N Autumn nitrogen fertiliser (kg N):

0	0
62	62

Pairs of sixth plots

2. N RATE Rate of nitrogen fertiliser (kg N):

50	50
100	100
150	150

NOTES: (1) AUT N 0 plots received a dressing of (0:20:20) at 160 kg.  
All other plots received a dressing of (20:10:10) at 310 kg.  
All broadcast at drilling.  
(2) Chlormequat was applied on 4 May.  
(3) N was applied to N RATE plots on 21 Apr.

Basal applications: Manures: (0:20:20) at 1260 kg applied to stubble before ploughing. Weedkillers: Methabenzthiazuron at 3.1 kg in 340 l in autumn. Isoproturon at 2.2 kg with mecoprop at 0.63 kg plus ioxynil at 0.21 kg in 340 l in spring.

Cultivations, etc.: - Basal PK applied: 29 Sept, 1975. Seed sown: 14 Oct.  
Autumn weedkiller applied: 15 Oct. Spring weedkillers applied: 25 Mar, 1976. Combine harvested: 27 July.

76/S/CS/1

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PREVCROP	7 WHEAT	8 WHEAT	9 WHEAT	10 WHEAT	MEAN
CCC					
0.0	4.49	4.66	4.32	4.67	4.54
1.7	5.08	4.73	4.88	4.44	4.78
N TIME					
SINGLE	4.82	4.69	4.60	4.59	4.68
DIVIDED	4.75	4.70	4.61	4.51	4.64
VARIETY					
CAPPELLE	4.64	4.61	4.53	4.43	4.56
HUNTSMAN	4.93	4.78	4.67	4.68	4.76
N RATE					
50	3.98	4.04	3.85	4.01	3.97
100	4.68	4.97	4.61	4.42	4.67
150	5.69	5.09	5.35	5.23	5.34
MEAN	4.79	4.70	4.60	4.55	4.66

GRAIN MEAN DM% 82.9

SUB PLOT AREA HARVESTED 0.00355

EXTRA

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

N RATE	50	100	150	MEAN
AUT N				
0	2.93	3.52	4.04	3.50
62	3.56	4.32	4.82	4.23
MEAN	3.24	3.92	4.43	3.87

GRAIN MEAN DM% 82.2

STRAW TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

N RATE	50	100	150	MEAN
AUT N				
0	2.18	2.59	3.20	2.65
62	3.03	3.68	4.32	3.68
MEAN	2.60	3.14	3.76	3.17

STRAW MEAN DM% 82.3

SUBPLOT AREA HARVESTED 0.00355