

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1976

[Full Table of Content](#)

76/R/WW/7 Rates and Times of N and K - W. Wheat

Rothamsted Research

Rothamsted Research (1977) *76/R/WW/7 Rates and Times of N and K - W. Wheat* ; Yields Of The Field Experiments 1976, pp 272 - 275 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/R/WW/7

WINTER WHEAT

RATES AND TIMES OF N AND K

Object: To study the effects of a range of rates and times of applying nitrogen and potassium on nutrient uptake and yield of winter wheat - Gt. Knott III.

Sponsors: O. Talibudeen, A. Penny, M. Page.

Design: Single replicate of 4 x 4 x 2 x 3 fully randomised plus 3 randomised blocks of 8 plots.

Whole plot dimensions: 2.67 x 4.57.

Treatments: All combinations of:-

1. N METHOD Rates, forms and times of applying nitrogen (kg N):

	'Nitro-Chalk' at G.S.2 (9 Apr)		Urea spray at G.S.6 (11-13 May)		Urea spray at G.S.11 (28-30 June)
120 0 30	120	+	0	+	30
120 0 60	120	+	0	+	60
120 0 90	120	+	0	+	90
120 30 0	120	+	30	+	0
120 30 30	120	+	30	+	30
120 30 60	120	+	30	+	60
120 30 90	120	+	30	+	90
120 60 0	120	+	60	+	0
120 60 30	120	+	60	+	30
120 60 60	120	+	60	+	60
120 60 90	120	+	60	+	90
120 90 0	120	+	90	+	0
120 90 30	120	+	90	+	30
120 90 60	120	+	90	+	60
120 90 90	120	+	90	+	90

2. N CONC Concentration of nitrogen in urea spray:-

2.5	2.5%
5.0	5.0%

3. K CONC Concentration of potassium in urea spray, K:N atom ratio:

0	None
5	5%
10	10%

EXTRA plus nine extra treatments given 'Nitro-Chalk' only, in spring at Growth Stage 2 (kg N) (9 Apr):

0	None
30	30
60	60
90	90
120 (duplicated)	120
150	150
180	180
240	240
300	300

76/R/WW/7

NOTE: Urea sprays were applied in 600 l to provide 30 kg N at N CONC 5.0 and in 1200 l to provide N CONC 2.5. The larger rates of urea were obtained by proportionate increases in volume.

Basal applications: Manures: (0:20:20) at 310 kg, combine drilled.

Weedkillers: Paraquat at 0.42 kg ion in 220 l. Ioxynil at 0.53 kg with mecoprop at 1.6 kg (both as the potassium salt) in 220 l in spring.

Fungicides: Tridemorph at 0.53 kg in 280 l. Benomyl at 0.28 kg with mancozeb plus maneb ('Kascade' at 2.2 kg) in 280 l. Insecticide: Pirimicarb at 0.14 kg in 280 l.

Seed: Cappelle, sown at 200 kg.

Cultivations, etc.:- Heavy spring-tine cultivated: 24 Sept, 1975. Paraquat applied: 6 Oct. Ploughed and rotary harrowed: 13 Oct. Seed sown: 14 Oct. Spring weedkiller applied: 22 Apr, 1976. Tridemorph applied: 29 Apr, 11 May. Benomyl with 'Kascade' applied, pirimicarb applied: 21 June. Combine harvested: 22 July. Previous crops: Barley 1974, winter oats 1975.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

N CONC	2.5	5.0	MEAN
N METHOD			
120 0 30	4.90	4.76	4.83
120 0 60	4.70	4.73	4.71
120 0 90	4.88	4.66	4.77
120 30 0	4.88	4.77	4.83
120 30 30	5.09	4.98	5.03
120 30 60	4.86	5.06	4.96
120 30 90	4.86	4.91	4.88
120 60 0	5.11	5.15	5.13
120 60 30	5.13	5.02	5.07
120 60 60	5.28	4.74	5.01
120 60 90	5.07	5.09	5.08
120 90 0	5.24	5.15	5.19
120 90 30	5.00	4.96	4.98
120 90 60	5.20	5.01	5.11
120 90 90	4.87	5.13	5.00
MEAN	5.00	4.94	4.97

76/R/WW/7

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

K CONC	0	5	10	MEAN
N METHOD				
120 0 30	4.53	4.79	5.10	4.83
120 0 60	4.53	4.75	4.86	4.71
120 0 90	4.86	4.87	4.53	4.77
120 30 0	4.88	4.91	4.69	4.83
120 30 30	5.14	5.12	4.84	5.03
120 30 60	5.05	5.04	4.78	4.96
120 30 90	4.86	4.93	4.81	4.88
120 60 0	5.26	5.11	5.01	5.13
120 60 30	4.90	5.03	5.24	5.07
120 60 60	5.09	5.09	4.85	5.01
120 60 90	5.13	4.96	5.15	5.08
120 90 0	5.48	4.88	5.22	5.19
120 90 30	4.65	5.25	5.03	4.98
120 90 60	4.94	5.36	5.01	5.11
120 90 90	5.15	4.86	4.97	5.00

MEAN 4.97 5.00 4.94 4.97

K CONC	0	5	10	MEAN
N CONC				
2.5	5.02	5.03	4.96	5.00
5.0	4.92	4.93	4.92	4.94

MEAN 4.97 5.00 4.94 4.97

EXTRA	0	30	60	90	120	150	180	240	300	MEAN
	3.25	3.45	4.30	4.47	4.89	5.22	4.79	5.29	5.06	4.56

GRAND MEAN 4.87

TABLE	N METHOD	N CONC	K CONC	EXTRA
SED	0.152	0.056	0.068	0.215 MIN REP 0.187 MAX-MIN
TABLE	N METHOD N CONC	N METHOD K CONC	N CONC K CONC	
SED	0.215	0.263	0.096	

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	47	0.264	5.4

EXTRA
MAX-MIN 120 V ANY OF REMAINDER
MIN REP ANY OF REMAINDER

GRAIN MEAN DM% 82.5

76/R/WW/7

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

N CONC	2.5	5.0	MEAN
N METHOD			
120 0 30	5.99	5.92	5.95
120 0 60	5.48	5.62	5.55
120 0 90	6.09	5.85	5.97
120 30 0	5.56	6.04	5.80
120 30 30	6.08	6.07	6.08
120 30 60	5.98	6.05	6.01
120 30 90	5.88	5.75	5.82
120 60 0	6.07	6.31	6.19
120 60 30	6.01	6.12	6.07
120 60 60	6.07	5.82	5.95
120 60 90	6.36	6.35	6.35
120 90 0	6.52	6.19	6.35
120 90 30	6.20	5.96	6.08
120 90 60	6.23	5.74	5.99
120 90 90	6.10	5.96	6.03

MEAN 6.04 5.98 6.01

K CONC	0	5	10	MEAN
N METHOD				
120 0 30	5.51	6.14	6.21	5.95
120 0 60	5.30	5.84	5.51	5.55
120 0 90	6.00	6.58	5.33	5.97
120 30 0	5.91	5.78	5.71	5.80
120 30 30	6.04	6.24	5.96	6.08
120 30 60	6.12	5.99	5.93	6.01
120 30 90	5.87	5.52	6.06	5.82
120 60 0	6.49	6.12	5.96	6.19
120 60 30	6.24	5.99	5.98	6.07
120 60 60	6.16	5.85	5.83	5.95
120 60 90	6.54	6.24	6.27	6.35
120 90 0	6.30	6.14	6.62	6.35
120 90 30	5.58	6.35	6.31	6.08
120 90 60	5.72	5.98	6.26	5.99
120 90 90	6.19	5.64	6.27	6.03

MEAN 6.00 6.03 6.01 6.01

K CONC	0	5	10	MEAN
N CONC				
2.5	5.95	6.09	6.09	6.04
5.0	6.05	5.97	5.94	5.98

MEAN 6.00 6.03 6.01 6.01

EXTRA	0	30	60	90	120	150	180	240	300	MEAN
	4.05	4.50	4.89	5.40	5.99	5.92	6.40	6.21	6.99	5.63

GRAND MEAN 5.92

STRAW MEAN DM% 89.6

PLOT AREA HARVESTED 0.00073