

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1976

[Full Table of Content](#)

### 76/R/WW/8 Fungicides and Grain Microflora - W. Wheat

#### Rothamsted Research

Rothamsted Research (1977) *76/R/WW/8 Fungicides and Grain Microflora - W. Wheat* ; Yields Of The Field Experiments 1976, pp 276 - 277 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/R/WW/8

WINTER WHEAT

FUNGICIDES AND GRAIN MICROFLORA

Object: To study the effects of a range of fungicides applied at a range of times on the yield, quality and grain microflora of winter wheat - Long Hoos IV 4.

Sponsor: R.A. Hill.

Design: Single replicate of 2 x 3 x 2 x 2 x 2 fully randomised.

Whole plot dimensions: 2.41 x 8.23.

Treatments: All combinations of:-

1. SPECFUNG Specific fungicides for foliar pathogen control:

NONE	None
T+B	Tridemorph at 0.53 kg plus benodanil at 1.4 kg mixed and applied on 14 and 27 May

2. B S FUNG Broad spectrum fungicides:

BENOMYL	Benomyl at 0.28 kg
CAPTAFOL	Captafol at 1.4 kg
BEN+CAP	Benomyl and captafol at above rates applied separately

3. APP TIME Application of broad spectrum fungicides:

	3 June	24 June	8 July
NONE	None	None	None
E	Sprayed	None	None
M	None	Sprayed	None
L	None	None	Sprayed
E+M	Sprayed	Sprayed	None
E+L	Sprayed	None	Sprayed
M+L	None	Sprayed	Sprayed
E+M+L	Sprayed	Sprayed	Sprayed

NOTE: All sprays were applied in 340 l.

Basal applications: Manures: (0:14:28) at 880 kg. 'Nitro-Chalk' at 450 kg.

Seed: Cama, sown at 200 kg.

Cultivations, etc.:- PK applied: 23 Sept, 1975. Ploughed: 26 Sept. Power harrowed and sown: 14 Oct. N applied: 10 May, 1976. Combine harvested: 28 July. Previous crops: Spring wheat 1974, fallow 1975.

NOTE: Grain microflora were assessed at fortnightly intervals from late May to harvest. Mildew and rust were assessed on several occasions. Eyespot and glume blotch were assessed before harvest.

76/R/WW/8

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

ALL PLOTS

APP TIME	NONE	E	M	L	E+M	E+L	M+L	E+M+L	MEAN
SPECFUNG									
NONE	5.84	5.83	5.89	5.93	5.84	6.03	5.54	5.58	5.82
T+B	5.68	5.93	6.19	5.72	5.65	6.00	5.87	5.87	5.86
MEAN	5.76	5.88	6.04	5.82	5.75	6.04	5.70	5.73	5.84

EXCLUDING APP TIME NONE

APP TIME	E	M	L	E+M	E+L	M+L	E+M+L	MEAN
B S FUNG								
BENOMYL	5.76	6.21	5.99	5.71	6.30	5.53	5.68	5.88
CAPTAFOL	5.98	5.60	6.26	5.54	5.83	5.73	5.57	5.79
BEN+CAP	5.90	6.31	5.21	5.98	6.01	5.85	5.93	5.89
MEAN	5.88	6.04	5.82	5.75	6.04	5.70	5.73	5.85

B S FUNG	BENOMYL	CAPTAFOL	BEN+CAP	MEAN
SPECFUNG				
NONE	5.76	5.69	5.98	5.81
T+B	6.00	5.88	5.79	5.89
MEAN	5.88	5.79	5.89	5.85

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	SPECFUNG	B S FUNG	APP TIME	SPECFUNG B S FUNG
SED	0.130 0.139*	0.170	0.260	0.241

TABLE	SPECFUNG APP TIME	B S FUNG APP TIME
SED	0.368	0.451

\* USE ONLY WITH TABLES EXCLUDING APP TIME NONE

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
WP	16	0.451	7.7

GRAIN MEAN DM% 87.8

PLOT AREA HARVESTED 0.00111