

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1976

[Full Table of Content](#)

A small thumbnail image of the document cover, which is a grid of numerical data. The columns are labeled with names like "SILVERHORN", "CLASSIFICATION", and "EXPERIMENTAL AREA". The rows contain various numbers, likely representing yield data for different treatments or plots.

76/R/WW/14 Seed Rates and Divided N Dressings - W. Wheat

Rothamsted Research

Rothamsted Research (1977) *76/R/WW/14 Seed Rates and Divided N Dressings - W. Wheat ; Yields Of The Field Experiments 1976*, pp 278 - 283 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/R/WW/14

WINTER WHEAT

SEED RATES AND DIVIDED N DRESSINGS

Object: To study the separate and combined effects of a range of treatments on the growth and yield of winter wheat - White Horse II.

Sponsors: G.V. Dyke, J. McEwen, R. Moffitt, D.B. Slope.

Design: Single replicate of 2 x 2 x 2 x 2 x 6 plus 4 extra plots.

Whole plot dimensions: 2.67 x 8.53.

Treatments: All combinations of:-

1. VARIETY Varieties:

CAPPELLE	Cappelle
HOBBIT	Hobbit

2. SEEDRATE Seed rates:

HALF	Half standard (100 kg Cappelle; 83 kg Hobbit) (1.86 million/ha)
STANDARD	Standard (200 kg Cappelle; 166 kg Hobbit) (3.72 million/ha)

3. CCC Chlormequat:

0.0	None
1.7	1.7 kg on 14 May

4. FUNGIDE Fungicide:

0	None
T+C+M	Tridemorph at 0.26 kg + carbendazim at 0.15 kg + maneb at 1.6 kg on 16 June

5. N Nitrogen fertiliser, times and rates (kg N):

	1 March	13 April	27 May
1 1 1	25	25	25
1 2 1	25	50	25
0 2 0	0	50	0
0 3 0	0	75	0
0 4 0	0	100	0
0 2 1	0	50	25

EXTRA plus four extra plots all sown at half standard seed rate, receiving chlormequat and the fungicidal mixture plus wettable sulphur at 2.66 kg a.i. in 340 l on 4 June:

C 111 S	Cappelle with N 1 1 1
C 121 S	Cappelle with N 1 2 1
H 111 S	Hobbit with N 1 1 1
H 121 S	Hobbit with N 1 2 1

76/R/WW/14

- NOTES: (1) All sprays were applied in 340 l.
(2) One plot VARIETY CAPPELLE, SEEDRATE STANDARD, CCC 0.0,
FUNGicide O, N 0 2 1 received CCC 1.7 in error. An
estimated value was used in the analysis.

Basal applications: Weedkillers: Dicamba with mecoprop and MCPA ('Banlene
Plus' at 5.6 l in 220 l).

Cultivations, etc.: - Deep-tine cultivated and heavy spring-tine cultivated:
7 Oct, 1975. Heavy spring-tine cultivated: 15 Oct. Rotary harrowed:
17 Oct. Heavy spring-tine cultivated: 23 Oct. Seed sown: 28 Oct.
Weedkiller applied: 20 Apr, 1976. Combine harvested: 3 Aug. Previous
crops: Beans 1974 and 1975.

NOTE: Plant counts were made in late autumn and spring. Samples were
taken for leaf area index, tiller numbers and dry weights in June.
Samples were taken for grain weights, straw weights, 1000 grain
weights and spikelet counts in late July.

76/R/WW/14

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

SEEDRATE VARIETY	HALFSTANDARD	MEAN					
CAPPELLE	3.99	4.75					
HOBBIT	3.99	4.77					
MEAN	3.99	4.76					
CCC VARIETY	0.0	1.7					
CAPPELLE	4.41	4.33					
HOBBIT	4.35	4.40					
MEAN	4.38	4.37					
CCC SEEDRATE	0.0	1.7					
HALF	4.05	3.93					
STANDARD	4.71	4.81					
MEAN	4.38	4.37					
FUNGicide VARIETY	0	T+C+M					
CAPPELLE	4.24	4.50					
HOBBIT	4.30	4.45					
MEAN	4.27	4.48					
FUNGicide SEEDRATE	0	T+C+M					
HALF	3.83	4.14					
STANDARD	4.71	4.81					
MEAN	4.27	4.48					
FUNGicide CCC	0.0	T+C+M					
1.7	4.30	4.47					
	4.25	4.48					
MEAN	4.27	4.48					
N VARIETY	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1	MEAN
CAPPELLE	4.23	4.27	4.59	4.37	4.30	4.49	4.37
HOBBIT	4.67	4.75	4.22	4.28	4.39	3.94	4.38
MEAN	4.45	4.51	4.41	4.32	4.34	4.22	4.37

76/R/WW/14

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1	MEAN
SEEDRATE								
HALF		4.21	3.96	3.84	3.90	4.16	3.86	3.99
STANDARD		4.69	5.06	4.97	4.74	4.53	4.57	4.76
MEAN		4.45	4.51	4.41	4.32	4.34	4.22	4.37
CCC	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1	MEAN
0.0		4.50	4.41	4.48	4.42	4.22	4.26	4.38
0.7		4.41	4.60	4.34	4.23	4.47	4.17	4.37
MEAN		4.45	4.51	4.41	4.32	4.34	4.22	4.37
FUNGICIDE	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1	MEAN
0		4.31	4.39	4.62	4.20	4.23	3.90	4.27
T+C+M		4.59	4.62	4.20	4.45	4.46	4.53	4.48
MEAN		4.45	4.51	4.41	4.32	4.34	4.22	4.37
SEEDRATE		HALF		STANDARD				
CCC		0.0		1.7		0.0		1.7
VARIETY	CAPPELLE		3.99		3.99		4.84	
	HOBBIT		4.10		3.87		4.59	
SEEDRATE		HALF		STANDARD				
FUNGICIDE		0		T+C+M		0		T+C+M
VARIETY	CAPPELLE		3.75		4.23		4.73	
	HOBBIT		3.92		4.05		4.69	
CCC		0.0			1.7			
FUNGICIDE		0		T+C+M		0		T+C+M
VARIETY	CAPPELLE		4.40		4.43		4.08	
	HOBBIT		4.19		4.51		4.42	
CCC		0.0			1.7			
FUNGICIDE		0		T+C+M		0		T+C+M
SEEDRATE		HALF						
		3.93		4.16		3.74		4.12
		STANDARD		4.77		4.77		4.85

76/R/WW/14

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1		
VARIETY	SEEDRATE								
CAPPELLE	HALF	3.93	3.67	4.22	3.80	4.09	4.24		
	STANDARD	4.53	4.86	4.97	4.93	4.50	4.74		
HOBBIT	HALF	4.49	4.24	3.47	4.01	4.22	3.49		
	STANDARD	4.86	5.25	4.97	4.55	4.56	4.40		
VARIETY	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1		
CAPPELLE	CCC								
	0.0	4.44	4.24	4.61	4.48	4.09	4.63		
	1.7	4.02	4.29	4.58	4.26	4.50	4.35		
HOBBIT	0.0	4.55	4.58	4.35	4.36	4.35	3.89		
	1.7	4.80	4.91	4.09	4.20	4.43	4.00		
SEEDRATE	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1		
HALF	CCC								
	0.0	4.23	3.90	3.92	4.21	3.96	4.07		
	1.7	4.19	4.01	3.77	3.59	4.35	3.66		
STANDARD	0.0	4.76	4.93	5.04	4.62	4.48	4.45		
	1.7	4.62	5.19	4.90	4.86	4.58	4.69		
VARIETY	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1		
CAPPELLE	FUNGicide								
	0	4.29	3.98	4.79	4.18	4.19	4.03		
	T+C+M	4.16	4.55	4.39	4.55	4.40	4.95		
HOBBIT	0	4.33	4.80	4.44	4.22	4.26	3.78		
	T+C+M	5.02	4.70	4.00	4.34	4.52	4.11		
SEEDRATE	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1		
FUNGicide									
HALF	0	4.14	3.82	4.09	3.91	3.81	3.24		
	T+C+M	4.28	4.09	3.60	3.89	4.51	4.49		
STANDARD	0	4.48	4.96	5.15	4.48	4.65	4.56		
	T+C+M	4.91	5.16	4.79	5.00	4.41	4.58		
CCC	N	1 1 1	1 2 1	0 2 0	0 3 0	0 4 0	0 2 1		
FUNGicide									
0.0	0	4.41	4.47	4.53	4.49	4.09	3.78		
	T+C+M	4.58	4.35	4.43	4.34	4.36	4.75		
1.7	0	4.21	4.30	4.71	3.90	4.36	4.03		
	T+C+M	4.60	4.90	3.96	4.55	4.57	4.32		
EXTRA	C 111 S	4.50	C 121 S	4.09	H 111 S	3.66	H 121 S	4.32	MEAN 4.15

76/R/WW/14

GRAIN TONNES/HECTARE

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	SEEDRATE	CCC	FUNGCIDE
SED	0.134	0.134	0.134	0.134
TABLE	N	VARIETY SEEDRATE	VARIETY CCC	SEEDRATE CCC
SED	0.233	0.190	0.190	0.190
TABLE	VARIETY FUNGCIDE	SEEDRATE FUNGCIDE	CCC FUNGCIDE	VARIETY N
SED	0.190	0.190	0.190	0.329
TABLE	SEEDRATE N	CCC N	FUNGCIDE N	VARIETY SEEDRATE CCC
SED	0.329	0.329	0.329	0.269
TABLE	VARIETY SEEDRATE FUNGCIDE	VARIETY CCC FUNGCIDE	SEEDRATE CCC FUNGCIDE	VARIETY SEEDRATE N
SED	0.269	0.269	0.269	0.466
TABLE	VARIETY CCC N	SEEDRATE CCC N	VARIETY FUNGCIDE N	SEEDRATE FUNGCIDE N
SED	0.466	0.466	0.466	0.466
TABLE	CCC FUNGCIDE N			
SED	0.466			

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	25	0.658	15.1

GRAIN MEAN DM% 37.3

PLOT AREA HARVESTED 0.00098