

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1976

[Full Table of Content](#)

### 76/S/WW/1 Rates and Times of N and Fungicide - W. Wheat

#### Rothamsted Research

Rothamsted Research (1977) *76/S/WW/1 Rates and Times of N and Fungicide - W. Wheat* ; Yields Of The Field Experiments 1976, pp 289 - 291 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/S/WW/1

WINTER WHEAT

RATES AND TIMES OF N AND FUNGICIDE

Object: To study the effects of rates and times of solid and liquid nitrogen fertilisers and foliar pathogen control on foliar disease incidence and yield of winter wheat - Saxmundham, Grove Plot.

Sponsors: F.V. Widdowson, A. Penny.

Design: Single replicate of 4 x 2 x 2 x 2 fully randomised.

Whole plot dimensions: 6.10 x 3.04.

Treatments: All combinations of:-

1. S N RATE Rates of solid nitrogen fertiliser ('Nitro-Chalk') (kg N):

0	None
50	50
100	100
150	150

2. S N TIME Times of applying solid nitrogen fertiliser:

APRIL	21 April
MAY	18 May

3. L N RATE Rates of liquid nitrogen fertiliser ('Agsol N26' - Urea + NH<sub>4</sub> NO<sub>3</sub>) (kg N):

0	None
25+25	25 on 15 June + 25 on 8 July

4. FUNGICIDE Fungicides:

NONE	None
BE+MZ+MB	Benomyl + mancozeb + maneb on 18 May and 15 June

NOTE: Benomyl applied at 0.28 kg with mancozeb plus maneb ('Kascade' at 2.24 kg) in 380 l.

Basal applications: Manures: (20:10:10) at 310 kg. Weedkillers: Methabenzthiazuron at 3.1 kg in 380 l in autumn. Isoproturon at 2.2 kg with mecoprop at 0.63 kg plus ioxynil at 0.21 kg in 340-l in spring. Growth regulator: Chlormequat at 1.7 kg in 340 l.

Seed: Maris Huntsman, sown at 190 kg.

Cultivations, etc.: - Ploughed: 2 Oct, 1975. Seed sown and fertiliser applied: 14 Oct. Autumn weedkiller applied: 15 Oct. Spring weedkiller applied: 25 Mar, 1976. Growth regulator applied: 4 May. Combine harvested: 20 July. Previous crops: Beans 1975, barley 1974.

76/S/WW/1

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

ALL PLOTS

S N RATE	0	50	100	150	MEAN
L N RATE					
0	6.15	6.46	6.68	6.86	6.54
25+25	6.27	6.69	6.36	6.73	6.51
MEAN	6.21	6.58	6.52	6.80	6.53

S N RATE	0	50	100	150	MEAN
FUNGICIDE					
NONE	6.33	6.52	6.60	6.61	6.51
BE+MZ+MB	6.10	6.63	6.45	6.99	6.54
MEAN	6.21	6.58	6.52	6.80	6.53

FUNGICIDE	NONE	BE+MZ+MB	MEAN
L N RATE			
0	6.63	6.45	6.54
25+25	6.39	6.64	6.51
MEAN	6.51	6.54	6.53

EXCLUDING S N RATE 0

S N RATE	50	100	150	MEAN
S N TIME				
APRIL	6.89	6.45	7.17	6.84
MAY	6.26	6.59	6.43	6.43
MEAN	6.58	6.52	6.80	6.63

L N RATE	0	25+25	MEAN
S N TIME			
APRIL	6.78	6.89	6.84
MAY	6.56	6.30	6.43
MEAN	6.67	6.59	6.63

FUNGICIDE	NONE	BE+MZ+MB	MEAN
S N TIME			
APRIL	6.61	7.06	6.84
MAY	6.54	6.32	6.43
MEAN	6.57	6.69	6.63

76/S/WW/1

GRAIN TONNES/HECTARE

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	L N RATE	FUNGCIDE	S N RATE	S N TIME
SED	0.138 0.160*	0.138 0.160*	0.196	0.160

TABLE	L N RATE FUNGCIDE	S N RATE S N TIME	S N RATE L N RATE	S N TIME L N RATE
SED	0.196	0.276	0.276	0.226

TABLE	S N RATE FUNGCIDE	S N TIME FUNGCIDE
SED	0.276	0.226

\* USE ONLY WITH TABLES EXCLUDING S N RATE 0

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
WP	13	0.391	6.0

GRAIN MEAN DM% 78.8

PLOT AREA HARVESTED 0.00089