

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1976

[Full Table of Content](#)

76/S/B/1 Varieties, N and Fungicides - Barley

Rothamsted Research

Rothamsted Research (1977) *76/S/B/1 Varieties, N and Fungicides - Barley* ; Yields Of The Field Experiments 1976, pp 317 - 320 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/S/B/1

SPRING BARLEY

VARIETIES, N AND FUNGICIDES

Object: To study the effects of rates, times and forms of nitrogen and of fungicides on the incidence of foliar diseases and yields of two varieties of barley - Saxmundham, Grove Plot.

Sponsors: F.V. Widdowson, J.F. Jenkyn, A. Penny.

Design: Single replicate of 32 plots split into 2.

Whole plot dimensions: 2.43 x 12.2.

Treatments: All combinations of:-

Whole plots

1. VARIETY Varieties:

JULIA	Julia
WING	Wing

2. S N Solid nitrogen fertiliser (kg N):

50	50
100	100

3. S N TIME Time of applying solid nitrogen fertiliser:

SEEDBED	Seedbed on 16 Mar
TOPDRESS	Top dressed on 18 May

4. L N Liquid nitrogen fertiliser (kg N):

0	None
50	50, half on 15 June, half on 8 July

5. MILD CON Mildew control:

NONE	None
ETH/TRID	Seed dressed eithirimol, crop sprayed tridemorph at 0.53 kg in 280 l on 18 May

Sub plots

6. RUST CON Rust control:

NONE	None
BENODANI	Crop sprayed benodanil at 1.12 kg in 280 l on 15 June, 8 July

NOTE: 'Nitro-Chalk' was used as solid fertiliser, 'Agsol N26' as liquid fertiliser in 75.

76/S/B/1

Basal applications: Manures: (0:20:20) at 315 kg. Weedkillers: Dichlorprop plus MCPA ('Mephetol Plus' at 8.4 l in 340 l).

Seed: Varieties sown at 190 kg.

Cultivations, etc.: - PK applied: 29 Sept, 1975. Ploughed: 20 Oct. Seed sown: 15 Mar, 1976. Weedkillers applied: 18 May. Combine harvested: 19 July. Previous crops: Barley 1974 and 1975.

NOTE: Brown rust and mildew were assessed in early July.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

S N	50	100	MEAN
VARIETY			
JULIA	3.40	3.31	3.35
WING	3.44	3.59	3.51
MEAN	3.42	3.45	3.43
S N TIME	SEEDBED	TOPDRESS	MEAN
VARIETY			
JULIA	3.51	3.19	3.35
WING	3.42	3.61	3.51
MEAN	3.46	3.40	3.43
S N TIME	SEEDBED	TOPDRESS	MEAN
S N			
50	3.34	3.49	3.42
100	3.59	3.31	3.45
MEAN	3.46	3.40	3.43
L N	0	50	MEAN
VARIETY			
JULIA	3.19	3.51	3.35
WING	3.40	3.63	3.51
MEAN	3.29	3.57	3.43
L N	0	50	MEAN
S N			
50	3.28	3.55	3.42
100	3.30	3.59	3.45
MEAN	3.29	3.57	3.43
L N	0	50	MEAN
S N TIME			
SEEDBED	3.14	3.78	3.46
TOPDRESS	3.44	3.36	3.40
MEAN	3.29	3.57	3.43

76/S/B/1

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

MILD CON	NONE	ETH/TRID	MEAN
VARIETY			
JULIA	3.48	3.22	3.35
WING	3.55	3.48	3.51
MEAN	3.51	3.35	3.43
MILD CON	NONE	ETH/TRID	MEAN
S N			
50	3.43	3.40	3.42
100	3.60	3.30	3.45
MEAN	3.51	3.35	3.43
MILD CON	NONE	ETH/TRID	MEAN
S N TIME			
SEEDBED	3.55	3.38	3.46
TOPDRESS	3.48	3.33	3.40
MEAN	3.51	3.35	3.43
MILD CON	NONE	ETH/TRID	MEAN
L N			
0	3.44	3.14	3.29
50	3.59	3.56	3.57
MEAN	3.51	3.35	3.43
RUST CON	NONE	BENODANI	MEAN
VARIETY			
JULIA	3.33	3.37	3.35
WING	3.44	3.58	3.51
MEAN	3.39	3.48	3.43
RUST CON	NONE	BENODANI	MEAN
S N			
50	3.43	3.40	3.42
100	3.34	3.56	3.45
MEAN	3.39	3.48	3.43
RUST CON	NONE	BENODANI	MEAN
S N TIME			
SEEDBED	3.41	3.51	3.46
TOPDRESS	3.36	3.44	3.40
MEAN	3.39	3.48	3.43
RUST CON	NONE	BENODANI	MEAN
L N			
0	3.26	3.32	3.29
50	3.51	3.63	3.57
MEAN	3.39	3.48	3.43

76/S/B/1

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

RUST CON	NONE	BENODANI	MEAN
MILD CON			
NONE	3.37	3.65	3.51
ETH/TRID	3.40	3.30	3.35
MEAN	3.39	3.48	3.43

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	VARIETY	S N	S N TIME	L N
SED	0.126	0.126	0.126	0.126

TABLE	MILD CON	RUST CON	VARIETY S N	VARIETY S N TIME
SED	0.126	0.113	0.179	0.179

TABLE	S N S N TIME	VARIETY L N	S N L N	S N TIME L N
SED	0.179	0.179	0.179	0.179

TABLE	VARIETY MILD CON	S N MILD CON	S N TIME MILD CON	L N MILD CON
SED	0.179	0.179	0.179	0.179

TABLE	VARIETY RUST CON	S N RUST CON	S N TIME RUST CON	L N RUST CON
SED	0.170	0.170	0.170	0.170

EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:
 VARIETY 0.160
 S N 0.160
 S N TIME 0.160
 L N 0.160

TABLE	MILD CON RUST CON
SED	0.170
EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:	
MILD CON	0.160

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	6	0.357	10.4
WP.SP	16	0.454	13.2

GRAIN MEAN DM% 84.0

SUB PLOT AREA HARVESTED 0.00089