


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1977

[Full Table of Content](#)



## 77/R/B/3 N and Foliar Diseases - Barley

### Rothamsted Research

Rothamsted Research (1978) 77/R/B/3 N and Foliar Diseases - Barley ; Yields Of The Field Experiments 1977, pp 346 - 349 - DOI: <https://doi.org/10.23637/ERADOC-1-29>

77/R/B/3  
SPRING BARLEY  
N AND FOLIAR DISEASES

Object: To study the effects of mildew and brown rust on response to a range of nitrogen rates applied at different times - Long Hoos I/II.

Sponsors: J.F. Jenkyn, M.E. Finney.

Design: Single replicate of 6 x 3 x 2 x 2.

Whole plot dimensions: 4.27 x 9.14.

Treatments: All combinations of:-

1. N RATE Amounts of nitrogen fertiliser (kg N):

25  
50  
70  
90  
110  
135

2. N TIME Times of applying N:

SB Seedbed (12 Apr, 1976)  
TD Top dressed (26 May)  
SB/TD Half to seedbed, half top dressed

3. MILDEW F Mildew fungicide:

NONE None  
TRIDEMOR Tridemorph on 20 June and 13 July

4. RUST F Rust fungicide:

NONE None  
BENODANI Benodanil on 11 July

NOTES: (1) Fungicides were applied in 340 l:-

(a) Tridemorph at 0.53 kg

(b) Benodanil at 1.12 kg with 175 ml 'Citowett'

(2) Sides of plots were separated by a strip of Mazurka barley 2.13 m wide.

Basal applications: Manures: (0:20:20) at 310 kg, combine drilled. Weedkillers: Dicamba with mecoprop and MCPA ('Banlene Plus' at 4.9 l in 220 l).

Seed: Zephyr, sown at 160 kg.

Cultivations, etc.: Ploughed: 30 Sept, 1976. Spring-tine cultivated, seed sown: 31 Mar, 1977. Weedkillers applied: 26 May. Combine harvested: 27 Aug.  
Previous crops: Potatoes 1975, wheat 1976.

NOTES: (1) Seedling emergence counts were made.

(2) Mildew and brown rust were assessed throughout the season.

(3) Samples were taken for N analysis.

(4) Ear counts were made before harvest.

77/R/B/3

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

N N TIME RATE	SB	TD	SB/TD	MEAN
25	4.99	5.00	5.00	4.99
50	5.18	5.54	5.60	5.44
70	5.48	5.49	5.71	5.56
90	5.54	5.66	6.09	5.76
110	5.73	5.48	6.05	5.75
135	5.77	5.75	6.08	5.86
MEAN	5.45	5.49	5.75	5.56
MILDEW F N RATE	NONE	TRIDEMOR	MEAN	
25	4.90	5.09	4.99	
50	5.34	5.55	5.44	
70	5.33	5.79	5.56	
90	5.49	6.03	5.76	
110	5.49	6.01	5.75	
135	5.59	6.14	5.86	
MEAN	5.36	5.77	5.56	
MILDEW F N TIME	NONE	TRIDEMOR	MEAN	
SB	5.30	5.59	5.45	
TD	5.23	5.75	5.49	
SB/TD	5.54	5.96	5.75	
MEAN	5.36	5.77	5.56	
RUST F N RATE	NONE	BENODANI	MEAN	
25	5.07	4.92	4.99	
50	5.56	5.33	5.44	
70	5.64	5.48	5.56	
90	5.73	5.79	5.76	
110	5.71	5.79	5.75	
135	5.76	5.96	5.86	
MEAN	5.58	5.55	5.56	
RUST F N TIME	NONE	BENODANI	MEAN	
SB	5.41	5.49	5.45	
TD	5.50	5.48	5.49	
SB/TD	5.83	5.68	5.75	
MEAN	5.58	5.55	5.56	

77/R/B/3

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

RUST F	NONE	BENODANI	MEAN			
MILDEW F						
NONE	5.39	5.33	5.36			
TRIDEMOR	5.77	5.77	5.77			
MEAN	5.58	5.55	5.56			
N TIME	SB		TD	SB/TD		
MILDEW F	NONE	TRIDEMOR	NONE	TRIDEMOR	NONE	TRIDEMOR
N RATE						
25	4.72	5.26	4.98	5.01	5.00	5.00
50	5.22	5.15	5.25	5.84	5.54	5.67
70	5.34	5.62	5.14	5.84	5.52	5.90
90	5.41	5.66	5.44	5.89	5.63	6.55
110	5.52	5.94	5.06	5.89	5.90	6.19
135	5.61	5.92	5.48	6.01	5.69	6.47
N TIME	SB		TD	SB/TD		
RUST F	NONE	BENODANI	NONE	BENODANI	NONE	BENODANI
N RATE						
25	5.16	4.82	5.21	4.79	4.85	5.15
50	5.17	5.20	5.69	5.40	5.82	5.39
70	5.55	5.41	5.57	5.41	5.79	5.63
90	5.35	5.72	5.56	5.76	6.28	5.90
110	5.67	5.78	5.33	5.63	6.12	5.97
135	5.54	5.99	5.62	5.87	6.13	6.03
MILDEW F	NONE		TRIDEMOR			
RUST F	NONE	BENODANI	NONE	BENODANI		
N RATE						
25	4.97	4.82	5.16	5.02		
50	5.48	5.19	5.64	5.47		
70	5.33	5.34	5.95	5.63		
90	5.56	5.43	5.91	6.16		
110	5.45	5.54	5.97	6.05		
135	5.54	5.65	5.99	6.28		
MILDEW F	NONE		TRIDEMOR			
RUST F	NONE	BENODANI	NONE	BENODANI		
N TIME						
SB	5.30	5.31	5.52	5.66		
TD	5.23	5.23	5.77	5.73		
SB/TD	5.64	5.45	6.02	5.91		

77/R/B/3

GRAIN TONNES/HECTARE

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	N RATE	N TIME	MILDEW F	RUST F
SED	0.093	0.066	0.054	0.054
TABLE	N RATE	N RATE	N TIME	N RATE
	N TIME	MILDEW F	MILDEW F	RUST F
SED	0.161	0.131	0.093	0.131
TABLE	N TIME	MILDEW F	N RATE	N RATE
	RUST F	RUST F	N TIME	N TIME
			MILDEW F	RUST F
SED	0.093	0.076	0.228	0.228
TABLE	N RATE	N TIME	N RATE	N RATE
	MILDEW F	MILDEW F	N TIME	N TIME
	RUST F	RUST F		
SED	0.186	0.131		

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
WP	10	0.228	4.1

GRAIN MEAN DM% 77.0

PLOT AREA HARVESTED 0.00195