


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1977

[Full Table of Content](#)



---

### 77/R/B/5 Mixed Varieties and Mildew - Barley

#### Rothamsted Research

Rothamsted Research (1978) *77/R/B/5 Mixed Varieties and Mildew - Barley* ; Yields Of The Field Experiments 1977, pp 354 - 355 - DOI: <https://doi.org/10.23637/ERADOC-1-29>

77/R/B/5

SPRING BARLEY

MIXED VARIETIES AND MILDEW

Object: To study mildew development and yield of variety mixtures sown at different seed rates, with and without a fungicidal seed dressing - Delafield.

Sponsor: J.F. Jenkyn.

Design: 3 randomised blocks of 12 plots.

Whole plot dimensions: 6.40 x 9.14.

Treatments: All combinations of:-

1. VAR SR            Varieties and seed rates:

HASSAN 2	Hassan, sown at 160 kg
MIDAS 2	Midas, sown at 160 kg
WING 2	Wing, sown at 160 kg
MIXED 1	Equal parts of the three varieties mixed and sown at 80 kg
MIXED 2	Equal parts of the three varieties mixed and sown at 160 kg
MIXED 3	Equal parts of the three varieties mixed and sown at 240 kg
  
2. FUNGICIDE        Systemic fungicidal seed dressing:

NONE	None
ETHIRIMO	Ethirimol at 15 g per 100 kg seed

NOTE: All plots were separated and surrounded by 12 m of variety Proctor, seed dressed ethirimol at 45 g per 100 kg seed, sprayed tridemorph at 0.53 kg in 340 l on 12 July, 1977. Yields were taken from this crop, adjacent to plots, and used for covariance analysis.

Basal applications: Manures: (20:14:14) at 380 kg, combine drilled. Weedkillers: Mecoprop with bromoxynil and ioxynil ('MAC CMPP' at 2.8 l with 'Oxytril CM' at 1.1 l in 220 l).

Cultivations, etc.: - Ploughed: 27 Oct, 1976. Spring-tine cultivated: 10 Mar, 1977. Sown: 12 Mar. Weedkillers applied: 24 May. Combine harvested: 5 Sept. Previous crops: Beans 1975, wheat 1976.

NOTE: Seedling emergence counts were made. Mildew was assessed on three occasions. Ear counts were made in August.

77/R/B/5

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

	VAR SR	HASSAN 2	MIDAS 2	WING 2	MIXED 1	MIXED 2	MIXED 3	MEAN
FUNGCIDE								
NONE		4.09	4.70	4.38	4.45	4.52	4.57	4.45
ETHIRIMO		3.92	5.03	4.44	4.40	4.42	4.78	4.50
MEAN		4.00	4.86	4.41	4.43	4.47	4.68	4.47

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	FUNGCIDE	VAR SR	FUNGCIDE VAR SR
SED	0.046	0.081	0.114

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	21	0.137	3.1

GRAIN MEAN DM% 83.6

PLOT AREA HARVESTED 0.00260