Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1977



Full Table of Content

77/R/RN/8 Cultivation/WEEDKILLER - S. Wheat

Rothamsted Research

Rothamsted Research (1978) 77/R/RN/8 Cultivation/WEEDKILLER - S. Wheat; Yields Of The Field Experiments 1977, pp 78 - 80 - DOI: https://doi.org/10.23637/ERADOC-1-29

77/R/RN/8

CULTIVATION/WEEDKILLER

Object: To study the long-term effects of weedkillers and different methods of primary cultivation on a rotation of crops - Great Harpenden I.

Sponsors: R. Moffitt, G.V. Dyke, J.A. Currie.

The 17th year, spring wheat.

For previous years see 'Details' 1967 and 1973 and 74-76/R/RN/8.

Design: 2 randomised blocks of 12 plots split into 2.

Whole plot dimensions: 12.8 x 15.2.

Treatments: All combinations of:-

Whole plots

1. CULTIVIN Primary cultivations annually:

Ploughed: 20 Oct, 1976 **PLOUGH** Rotary cultivated: 13 Oct ROTAVATE

Deep-tine cultivated twice: 5 Oct DEEPTINE

Weed control to beans 1976: 2. WEEDCNTL(76)

Mechanical MECHANCL SIMAZINE Simazine

Sub plots

Hormone weedkiller to previous cereals, last applied to WEEDKLLR(751) barley 1975 (basal weedkiller 'Banlene Plus' at 4.9 1

in 220 1 in 1977):

NONE None

DI+ME+MC Dicamba + mecoprop + MCPA

Paraquat weedkiller to previous cereal stubbles, last 4. WEEDKLLR(752)

applied to barley stubble autumn 1975 (basal weedkiller, glyphosate at 1.7 kg in 220 l to bean stubble 1976):

NONE PARAQUAT

EXTRA plus three extra whole plot treatments:

Heavy spring-tine cultivated twice: 6 Oct, 1976. Given SPNGTINE

simazine to beans 1976, with sub plot treatments 3 and

Shallow ploughed: 7 Oct, 1976. Given simazine to beans (SH)PLGH

1976 and paraquat to barley stubble autumn 1975, with sub

plot treatment 3 above.

Standard cultivations as considered best for each crop. STANDARD

Ploughed 20 Oct, 1976. Given simazine to beans 1976, with

sub plot treatments 3 and 4 above.

77/R/RN/8

NOTE: It was intended to sow winter wheat but prolonged wet weather prevented this.

Basal applications: Manures: (10:24:24) at 250 kg, combine drilled. Insecticide: Pirimicarb at 0.14 kg in 270 l.

Seed: Sappo, sown at 190 kg.

Cultivations, etc.:- Glyphosate applied: 6 Sept, 1976. All plots heavy springtine cultivated twice: 25 Nov, 16 Dec. All plots rotary harrowed: 16 Dec. All plots spring-tine cultivated: 4 Apr, 1977. Seed sown: 5 Apr. 'Banlene Plus' applied: 26 May. Insecticide applied: 14 July. Combine harvested: 23 Sept.

EXTRA PLOTS ONLY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

EXTRA WEEDKLLR(751)	SPNGTINE	(SH)PLGH	STANDARD
NONE D1+ME+ML	4.32 4.82	4.77 5.25	4.32 4.48
WEEDKLLR(752) NONE PARAQUAT	4.62 4.53	5.01	4.39 4.41
MEAN	4.58	5.01	4.40

GRAIN MEAN DM% 76.3

SUB PLOT AREA HARVESTED 0.00434

77/R/RN/8

OMITTING EXTRA PLOTS

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

CULTIVTN	PLOUGH	ROTAVATE	DEEPTINE	MEAN
WEEDCNTL(76) MECHANCL SIMAZINE	4.26 4.70	4.04 4.45	4.82 4.70	4.37 4.62
WEEDKLLR(751) NONE D1+ME+ML	4.62 4.49	4.33 4.30	4.71 4.77	4.55 4.52
WEEDKLLR(752) NONE PARAQUAT	4.56 4.55	4.31 4.33	4.78 4.70	4.55 4.53
MEAN	4.55	4.32	4.74	4.54

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CULTIVIN	WEEDCNTL(76)	WEEDKLLR(751)	WEEDKLLR(752)
SED	0.178	0.154	0.038	0.038
TABLE	CULTIVIN WEEDCNTL(76)	CULTIVIN WEEDKLLR(751)	CULTIVIN WEEDKLLR(752)	
SED	0.309 0.268 0.218	0.184	0.184	MIN REP MAX-MIN MAX REP

EXCEPT WHEN COMPARING MEANS WITH SAME LEVEL(S) OF:

CULTIVIN

0.066 0.066

WEEDCNTL(76)

MIN REP MECHANCL MAX-MIN MECHANCL V SIMAZINE MAX REP SIMAZINE

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	11	0.309	6.8
BLOCK.WP.SP	10	0.114	2.5

GRAIN MEAN DM% 76.4

SUB PLOT AREA HARVESTED 0.00434