


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1978

[Full Table of Content](#)



---

### 78/W/RN/13 Intensive Cereals - Wheat, Barley

#### Rothamsted Research

Rothamsted Research (1979) *78/W/RN/13 Intensive Cereals - Wheat, Barley* ; Yields Of The Field Experiments 1978, pp 103 - 106 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/W/RN/13

INTENSIVE CEREALS

Object: To study the effects of intensive cereal cropping on yield, incidence of soil-borne pathogens and organic matter in the soil - Woburn Stackyard I.

Sponsors: A.E. Johnston, J. McEwen.

The 13th year, winter wheat, barley.

For previous years see 'Details' 1973 and 74-77/W/RN/13.

Design: For each experiment: 2 randomised blocks of 6 plots, split into 4. ALDICARB tested on blocks.

Whole plot dimensions: 8.53 x 20.4.

Treatments:-

One experiment on winter wheat on part of the site of the classical wheat experiment 1877-1954

One experiment on barley on part of the site of the classical barley experiment 1877-1954

Factors tested on both experiments are the same but crop and nitrogen rates differ. All combinations of:-

Blocks

1. ALDICARB Aldicarb, cumulative to 1977 dressing, worked into the seedbed (kg):

0  
10

Whole plots

2. PREVCROP Previous crops:

	1972	1973	1974	1975	1976	1977
C/L/P/C	C	C	C	L	P	C
L/P/C/C	C	C	L	P	C	C
P/C/C/C	C	L	P	C	C	C
C/C/C/C	L	P	C	C	C	C
C/C/L/C	P	C	C	C	L	C
C/C/C/C	C	C	C	C	C	C

Ley = 1 year ley P = Potatoes C = Cereal: wheat or barley. All plots in cereal from 1977.

Sub plots

3. N Nitrogen fertiliser (kg N as 'Nitro-Chalk 25'):

Wheat	Barley
63	50
126	100
189	150
252	200

78/W/RN/13

Standard applications:

Wheat: Manures: (0:20:20) at 310 kg, combine drilled. Weedkillers: Methabenzthiazuron at 1.5 kg in 280 l. Mecoprop, bromoxynil and ioxynil ('Brittox' at 3.5 l in 280 l).

Barley: Manures: (0:20:20) at 300 kg, combine drilled. Weedkillers: Mecoprop, bromoxynil and ioxynil ('Brittox' at 2.5 l in 280 l). Fungicide: Tridemorph at 0.53 kg applied with the weedkillers.

Seed: Wheat: Cappelle, sown at 210 kg.

Barley: Porthos, dressed with ethirimol, sown at 160 kg.

Cultivations, etc.:— All plots ploughed: 27 Sept, 1977.

Wheat: Aldicarb applied, rotary cultivated: 24 Oct, 1977. Spring-tine cultivated, seed sown: 25 Oct. Methabenzthiazuron applied: 29 Oct. N applied: 7 Apr, 1978. Mecoprop, bromoxynil and ioxynil applied: 10 May. Combine harvested: 25 Aug.

Barley: Spring-tine cultivated: 9 Mar, 1978. Aldicarb applied, rotary cultivated, spring-tine cultivated with crumbler attached, seed sown: 3 Apr. N applied, weedkillers and fungicide applied: 15 May. Combine harvested: 23 Aug.

78/W/RN/13

WHEAT

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PREVCROP	C/L/P/C	L/P/C/C	P/C/C/C	C/C/C/C	C/C/L/C	C/C/C/C	MEAN
ALDICARB							
0	4.26	4.26	3.88	3.81	4.36	3.94	4.08
10	4.06	3.92	3.74	3.44	3.62	3.18	3.66
MEAN	4.16	4.09	3.81	3.62	3.99	3.56	3.87
N	63	126	189	252	MEAN		
ALDICARB							
0	3.42	4.32	4.36	4.23	4.08		
10	3.15	3.95	3.66	3.87	3.66		
MEAN	3.29	4.14	4.01	4.05	3.87		
N	63	126	189	252	MEAN		
PREVCROP							
C/L/P/C	3.52	4.71	4.39	4.01	4.16		
L/P/C/C	3.76	4.22	4.01	4.38	4.09		
P/C/C/C	3.21	3.98	4.06	3.99	3.81		
C/C/C/C	2.98	3.65	3.95	3.91	3.62		
C/C/L/C	3.35	4.33	3.84	4.43	3.99		
C/C/C/C	2.90	3.94	3.83	3.58	3.56		
MEAN	3.29	4.14	4.01	4.05	3.87		
N	63	126	189	252			
ALDICARB	PREVCROP						
0	C/L/P/C	3.45	4.56	4.51	4.51		
	L/P/C/C	3.92	4.79	4.49	3.83		
	P/C/C/C	2.83	3.98	3.98	4.73		
	C/C/C/C	3.47	3.70	4.29	3.77		
	C/C/L/C	3.29	4.73	4.73	4.70		
	C/C/C/C	3.57	4.18	4.18	3.83		
10	C/L/P/C	3.59	4.87	4.26	3.51		
	L/P/C/C	3.59	3.65	3.52	4.92		
	P/C/C/C	3.60	3.97	4.14	3.25		
	C/C/C/C	2.50	3.59	3.60	4.05		
	C/C/L/C	3.41	3.93	2.96	4.17		
	C/C/C/C	2.23	3.69	3.49	3.33		

GRAIN MEAN DM% 82.4

SUB PLOT AREA HARVESTED 0.00277

78/W/RN/13

BARLEY

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

PREVCROP	C/L/P/C	L/P/C/C	P/C/C/C	C/C/C/C	C/C/L/C	C/C/C/C	MEAN
ALDICARB							
0	5.00	5.00	4.63	4.35	4.07	3.81	4.48
10	4.75	4.94	4.65	3.95	4.80	4.20	4.55
MEAN	4.88	4.97	4.64	4.15	4.43	4.01	4.51
N	50	100	150	200	MEAN		
ALDICARB							
0	3.02	4.61	5.22	5.05	4.48		
10	2.87	4.74	5.17	5.42	4.55		
MEAN	2.94	4.67	5.20	5.24	4.51		
N	50	100	150	200	MEAN		
PREVCROP							
C/L/P/C	3.56	5.04	5.73	5.17	4.88		
L/P/C/C	3.82	5.06	5.28	5.72	4.97		
P/C/C/C	3.05	4.75	5.37	5.39	4.64		
C/C/C/C	1.97	4.59	5.18	4.87	4.15		
C/C/L/C	2.94	4.47	5.01	5.32	4.43		
C/C/C/C	2.34	4.13	4.62	4.94	4.01		
MEAN	2.94	4.67	5.20	5.24	4.51		
N	50	100	150	200			
ALDICARB							
PREVCROP							
0	C/L/P/C	4.06	4.89	6.06	5.00		
	L/P/C/C	3.85	5.28	5.46	5.39		
	P/C/C/C	3.30	4.46	5.21	5.54		
	C/C/C/C	2.39	4.76	5.23	5.03		
	C/C/L/C	2.49	4.21	4.85	4.72		
	C/C/C/C	2.05	4.08	4.51	4.62		
10	C/L/P/C	3.06	5.20	5.40	5.35		
	L/P/C/C	3.78	4.84	5.09	6.05		
	P/C/C/C	2.79	5.04	5.53	5.24		
	C/C/C/C	1.55	4.42	5.12	4.71		
	C/C/L/C	3.38	4.74	5.17	5.91		
	C/C/C/C	2.63	4.18	4.73	5.27		

GRAIN MEAN DM% 77.6

SUB PLOT AREA HARVESTED 0.00277