


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1978

[Full Table of Content](#)



78/W/CS/66 Dazomet and Nitrogen - Maize

Rothamsted Research

Rothamsted Research (1979) *78/W/CS/66 Dazomet and Nitrogen - Maize* ; Yields Of The Field Experiments 1978, pp 162 - 163 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/W/CS/66

DAZOMET AND NITROGEN

Object: To study the cumulative effects of dazomet and nitrogen on pathogens and yield of maize grown continuously - Woburn Butt Furlong.

Sponsors: A.J. Barnard, D. Hornby.

The eighth year, maize.

For previous years see 71/W/CS/66(t), 72/W/CS/66(t) and 73-77/W/CS/66.

Design: 4 blocks of 2 plots split into 4.

Whole plot dimensions: 2.13 x 16.5.

Treatments: All combinations of:-

Whole plots

1. DAZOMET Dazomet (kg per annum) cumulative 1971-78:

0
450

Sub plots

2. N Nitrogen fertiliser (kg N as 'Nitro-Chalk 34') cumulative 1971-78:

50 50 to seedbed
100 100 to seedbed
150 150 to seedbed
100+50 100 to seedbed, 50 five weeks before sowing (before 1978 this treatment received 100 to seedbed, 50 five weeks after emergence)

Basal applications: Manures: Magnesian limestone at 7.5 tonnes, (0:14:28) at 860 kg. Weedkiller: Atrazine at 1.1 kg in 340 l.

Seed: Fronica, sown at 103,300 seeds per ha.

Cultivations, etc.: - Magnesian limestone applied: 18 Nov, 1977. Ploughed: 21 Nov. Spring-tine cultivated with crumbler attached, early N applied, dazomet applied, rotary cultivated twice: 3 Apr, 1978. PK applied, spring-tine cultivated with crumbler attached, seed sown: 12 May. Seedbed N applied: 19 May. Weedkiller applied: 23 May. Hand harvested: 11 Oct.

NOTES: (1) Soil samples were taken before sowing and after harvest for counts of ectoparasitic nematodes.
(2) Counts were made of common smut (*Ustilago maydis*) and stalk rots (*Fusarium* spp.).

78/W/CS/66

FORAGE DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

	N	50	100	150	100+50	MEAN
DAZOMET						
0		8.56	9.89	11.26	11.49	10.30
450		10.61	11.99	13.51	13.70	12.45
MEAN		9.59	10.94	12.38	12.60	11.38

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	N	DAZOMET*
		N
SED	0.683	0.966

* WITHIN THE SAME LEVEL OF DAZOMET ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	18	1.366	12.0

FORAGE MEAN DM% 25.0

SUB PLOT AREA HARVESTED 0.00039