


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1978

[Full Table of Content](#)



78/W/CS/99 Effects of Breaks on Take-all - Barley, Beans

Rothamsted Research

Rothamsted Research (1979) *78/W/CS/99 Effects of Breaks on Take-all - Barley, Beans* ; Yields Of The Field Experiments 1978, pp 166 - 167 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/W/CS/99

EFFECTS OF BREAKS ON TAKE-ALL

Object: To study the phenomenon of take-all (*Gaeumannomyces graminis*) decline in barley - Woburn Butt Furlong.

Sponsor: D. Hornby.

The sixth year, barley, spring beans.

For previous years see 72/W/CS/99(t) and 73-77/W/CS/99.

Design: 2 blocks of 9 plots.

Whole plot dimensions: 5.33 x 15.2.

Treatments: Previous crops:-

PREVCROP	1968-71	1972	1973	1974	1975	1976	1977
BBBBBB	B	B	B	B	B	B	B
FBBBBB	B	F	B	B	B	B	B
FBEBBB	B	F	BE	B	B	B	B
BFEBBB	B	B	F	BE	B	B	B
BBFBEB	B	B	B	F	BE	B	B
BBBFEB	B	B	B	B	F	BE	B
BBBBFBE	B	B	B	B	B	F	BE

B = Barley, BE = Beans, F = Fallow

Standard applications:

All plots: Manures: Magnesian limestone at 7.5 tonnes.

Barley: Manures: (20:14:14) at 450 kg, combine drilled. Weedkillers: Mecoprop, bromoxynil and ioxynil ('Brittox' at 2.5 l in 280 l) applied with Fungicide: Tridemorph at 0.53 kg.

Spring beans: Manures: (0:14:28) at 400 kg. Insecticide: Pirimicarb at 0.14 kg in 280 l.

Seed: Barley: Porthos, dressed with ethirimol, sown at 160 kg.

Spring beans: Minden, sown at 220 kg.

Cultivations, etc.:-

All plots: Magnesian limestone applied: 17 Oct, 1977. Ploughed: 8 Nov. Spring-tine cultivated: 8 Mar, 1978.

Barley: Seed sown: 9 Mar, 1978. Weedkillers and fungicide applied: 15 May. Combine harvested: 18 Aug.

Spring beans: PK applied, seed sown: 15 Mar, 1978. Hand hoed twice: 24 May, 28 June. Insecticide applied: 14 June. Hand harvested: 11 Sept.

Fallow: Rotary cultivated: 16 May, 1978. Hand hoed twice: 24 May, 30 July.

- NOTES: (1) Soil samples were taken before sowing and after harvest and plant samples in July for incidence of take-all (*Gaeumannomyces graminis*).
(2) An examination of the results showed a fertility trend coinciding with the contours of the field. The yields presented have been adjusted for this trend.

78/W/CS/99

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVCROP	BBBBBB	FBBBBB	FBEBBBB	BFBE BBB	BBFBEBB	BBBFBE B	BBBBFBE	MEAN
	3.14	3.68	4.05	4.19	4.06	3.56	4.46	3.88

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	PREVCROP
-----	-----
SED	0.245

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	5	0.229	5.9

GRAIN MEAN DM% 83.5

PLOT AREA HARVESTED 0.00434