

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1978

[Full Table of Content](#)



78/R/CS/202 Effects of Phialophora - Wheat

Rothamsted Research

Rothamsted Research (1979) *78/R/CS/202 Effects of Phialophora - Wheat* ; Yields Of The Field Experiments 1978, pp 273 - 274 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/R/CS/202

EFFECTS OF PHIALOPHORA

Object: To study the effects of ryegrass, oats and wheat, and of soil inoculation on populations of *Phialophora radiculicola* (Prg) and on take-all (*Gaeumannomyces graminis*) and yield of following wheat crops - Whittlocks.

Sponsors: E. Lester, D.B. Slope, R.J. Gutteridge.

The second year, wheat.

For previous year see 77/R/CS/202.

Design: 3 randomised blocks of 8 plots.

Whole plot dimensions: 2.67 x 6.10.

Treatments:

CRP INOC(778) Crop in 1977 and inoculation in 1977 and 1978:

GRASS	Ryegrass
GRASS(I)	Ryegrass + Prg inoculum 1977
OATS	Spring oats
OATS(I)	Spring oats + Prg inoculum 1977
OATS I	Spring oats + Prg inoculum to 1978 wheat
OATS DI	Spring oats + dummy inoculum (sand) to 1978 wheat
WHEAT	Spring wheat
WHEAT(I)	Spring wheat + Prg inoculum 1977

NOTE: *Phialophora* inoculum was applied on 20 Oct, 1977. The inoculum was a *Phialophora* culture macerated in coarse sand. Coarse sand alone was applied as dummy inoculum on the same date.

Basal applications: Manures: (10:24:24) at 200 kg, combine drilled. 'Nitro-chalk 26' at 300 kg. K_2O at 100 kg as muriate of potash. Weedkillers: Paraquat at 0.56 kg ion² in 220 l. Terbutryne at 2.8 kg in 220 l.

Seed: Cappelle, sown at 190 kg.

Cultivations, etc.: Paraquat applied: 26 Sept, 1977. K applied, ploughed: 11 Oct. Rolled, rotary harrowed: 20 Oct. Power harrowed, seed sown and terbutryne applied: 24 Oct. N applied: 4 May, 1978. Combine harvested: 4 Sept, 1978.

NOTE: Take-all and *Phialophora* were estimated in plants in April and July and in soil in September.

78/R/CS/202

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

CRP INOC(778)	
GRASS	8.23
GRASS(I)	7.76
OATS	7.22
OATS(I)	7.82
OATS I	8.44
OATS DI	7.26
WHEAT	6.40
WHEAT(I)	5.75
MEAN	7.36

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CRP INOC(778)
-----	-----
SED	0.457

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	14	0.560	7.6

GRAIN MEAN DM% 80.6

PLOT AREA HARVESTED 0.00116