

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1978

[Full Table of Content](#)



---

### 78/R/CS/206 Late N - Wheat

#### Rothamsted Research

Rothamsted Research (1979) *78/R/CS/206 Late N - Wheat* ; Yields Of The Field Experiments 1978, pp 288 - 289 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/R/CS/206

LATE N

Object: To study the residual effects on wheat of a range of fertilisers applied to potatoes in 1977 - Great Knott II.

Sponsors: T.M. Addiscott, J. Ashworth, A. Penny, F.V. Widdowson.

The second year, wheat.

For first year see 77/R/CS/206.

Design: 3 randomised blocks of 18 plots.

Whole plot dimensions: 4.27 x 13.1.

Treatments: All combinations of:-

1. N FORM(77)      Forms of nitrogen fertiliser in 1977:
  - AA                    Aqueous ammonia, injected before planting
  - AA+NITRA            Aqueous ammonia + nitrapyrin ('N-Serve') at 1 kg, injected before planting
  - AA+STC              Aqueous ammonia + sodium trithiocarbonate at 42 kg injected before planting
  - IB SMALL            IBDU (isobutylidene diurea), small granules to seedbed
  - IB LARGE            IBDU, large granules to seedbed
  - AN E+L              Ammonium nitrate, half in seedbed, half top-dressed
  - AN E                 Ammonium nitrate, all in seedbed

2. N RATE(1)      Rates of nitrogen fertiliser in 1977 (kg N):

200  
300

plus four extra treatments given ammonium nitrate, all in the seedbed in 1977 (kg N):

N RATE(2)

AN E 150  
AN E 250  
AN E 350  
AN E 400

NOTES: (1) The intended rate of sodium trithiocarbonate was 24 kg. Only one plot, treatment N RATE(1)200, received this rate, all other plots received 42 kg.

(2) Aqueous ammonia and inhibitors were injected, 10 cm deep with tines 30 cm apart.

Basal applications: Manures: (0:20:20) at 310 kg, combine drilled. 'Nitra-Shell 34' at 240 kg. Weedkillers: Dicamba with mecoprop and MCPA ('Banlene Plus' at 4.9 l in 220 l). Growth regulator: Chlormequat at 1.4 kg in 220 l.

Seed: Atou, sown at 210 kg.

78/R/CS/206

Cultivations, etc.:— Heavy spring-tine cultivated twice: 7 and 8 Nov, 1977. Seed sown: 15 Nov. N applied: 19 Apr, 1978. Weedkiller and growth regulator applied: 10 May. Combine harvested: 12 Sept.

NOTE: Crop samples were taken for N determinations.

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

N RATE(1) N FORM(77)	200	300	MEAN
AA	5.01	5.69	5.35
AA+NITRA	5.26	5.89	5.57
AA+STC	5.41	5.96	5.69
IB SMALL	5.51	5.78	5.65
IB LARGE	5.63	5.90	5.76
AN E+L	5.91	5.99	5.95
AN E	5.33	5.79	5.56
MEAN	5.44	5.86	5.65

  

N RATE(2)	AN E 150	AN E 250	AN E 350	AN E 400	MEAN
	5.00	5.61	5.49	6.01	5.53

GRAND MEAN 5.62

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	N RATE(2)	N FORM(77)	N RATE(1)	N FORM(77) N RATE(1) & RATE(2)
SED	0.263	0.186	0.099	0.263

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	34	0.322	5.7
PLOT AREA HARVESTED	0.00279		