


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1978

[Full Table of Content](#)



## 78/S/WW/1 Rates and Times of N and Fungicides - W. Wheat

**Rothamsted Research**

Rothamsted Research (1979) *78/S/WW/1 Rates and Times of N and Fungicides - W. Wheat ; Yields Of The Field Experiments 1978*, pp 365 - 368 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/S/WW/1

WINTER WHEAT

RATES AND TIMES OF N AND FUNGICIDE

Object: To study the effects of fungicides and rates and times of nitrogen fertiliser on the incidence of foliar diseases and on yield of winter wheat - Saxmundham.

Sponsors: F.V. Widdowson, A. Penny.

Design: Half replicate of  $4 \times 2^4$  plus 8 extra plots.

Whole plot dimensions: 6.40 x 2.74.

Treatments: Combinations of:

1. N AUTUMN      Nitrogen fertiliser in autumn (19 Oct, 1977):

0	None
IBDU 1	Isobutylidene diurea at 50 kg N

2. N SPRING      Nitrogen fertiliser in spring (14 Mar, 1978):

0	None
NC 1	'Nitro-Chalk' 25% N at 50 kg N
NC 2	'Nitro-Chalk' 25% N at 100 kg N
NC 3	'Nitro-Chalk' 25% N at 150 kg N

3. N SUMMER      Nitrogen fertiliser in summer:

0	None
AG 1	'Agsol 26% N' at 50 kg N. Foliar spray, half on 7 June, half on 21 June

4. FUNGCIDE(1)    Fungicide:

0	None
BN+CA+MA	Benomyl on 18 May, carbendazim + maneb on 7 June and 6 July

5. FUNGCIDE(2)    Fungicide:

0	None
BENODANI	Benodanil on 21 June and on 6 July

plus four extra treatments (duplicated), all given FUNGCIDE(1) and FUNGCIDE(2):

EXTRA

NCA1NCD2	'Nitro-Chalk' in autumn at 50 kg N, 'Nitro-Chalk' in spring/summer at 100 kg N dressing divided 1/5 at G.S.3, 3/5 at G.S.5, 1/5 at G.S.8.
NCA1NCD3	As previous treatment but spring/summer dressing at 150 kg N
IBA1NCD2	Isobutylidene diurea in autumn at 50 kg N, 'Nitro-Chalk' in spring/summer at 100 kg N dressing divided as above
IBA1NCD3	As previous treatment but spring/summer dressing at 150 kg N

78/S/WW/1

- NOTES: (1) EXTRA nitrogen treatments were applied on the following dates:  
G.S.3 14 Mar, G.S.5 20 Apr, G.S.8 18 May.  
(2) 'FUNGicide(1)' Benomyl was applied at 0.28 kg in 280 l and  
carbendazim at 0.25 kg plus maneb at 1.6 kg in 280 l.  
(3) 'FUNGicide(2)' Benodanil was applied at 1.1 kg in 280 l.

Basal applications: Manures: Muriate of potash at 250 kg. (0:20:20) at 250 kg.  
Weedkillers: Isoproturon at 1.8 kg in 450 l. Ioxynil at 0.53 kg with  
mecoprop at 1.9 kg in 280 l applied with the fungicide and growth regulator  
(see below). Fungicide: Tridemorph at 0.53 kg. Insecticide: Pirimicarb at  
0.14 kg in 280 l. Growth regulator: Chlormequat at 1.7 kg.

Seed: Maris Huntsman, sown at 200 kg.

Cultivations, etc.: - Muriate of potash applied: 21 Sept, 1977. Ploughed: 24  
Sept. Harrowed and rolled three times: 12 Oct. PK applied, seed sown,  
isoproturon applied: 19 Oct. Ioxynil, mecoprop, tridemorph and chlormequat  
applied: 18 May, 1978. Pirimicarb applied: 3 Aug. Combine harvested:  
23 Aug.

78/S/WW/1

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

N SPRING	0	NC 1	NC 2	NC 3	MEAN
N AUTUMN					
0	2.83	5.20	6.37	6.88	5.32
IBDU 1	3.98	5.90	6.67	7.01	5.89
MEAN	3.40	5.55	6.52	6.94	5.60
N SUMMER	0	AG 1	MEAN		
N AUTUMN					
0	5.00	5.63	5.32		
IBDU 1	5.74	6.04	5.89		
MEAN	5.37	5.84	5.60		
N SUMMER	0	AG 1	MEAN		
N SPRING					
0	3.06	3.75	3.40		
NC 1	5.18	5.93	5.55		
NC 2	6.45	6.58	6.52		
NC 3	6.79	7.09	6.94		
MEAN	5.37	5.84	5.60		
FUNGCIDE(1)	0	BN+CA+MA	MEAN		
N AUTUMN					
0	5.23	5.41	5.32		
IBDU 1	5.80	5.98	5.89		
MEAN	5.51	5.69	5.60		
FUNGCIDE(1)	0	BN+CA+MA	MEAN		
N SPRING					
0	3.27	3.54	3.40		
NC 1	5.61	5.50	5.55		
NC 2	6.40	6.63	6.52		
NC 3	6.77	7.11	6.94		
MEAN	5.51	5.69	5.60		
FUNGCIDE(1)	0	BN+CA+MA	MEAN		
N SUMMER					
0	5.25	5.49	5.37		
AG 1	5.78	5.90	5.84		
MEAN	5.51	5.69	5.60		
FUNGCIDE(2)	0	BENODANI	MEAN		
N AUTUMN					
0	5.36	5.27	5.32		
IBDU 1	5.97	5.81	5.89		
MEAN	5.67	5.54	5.60		

78/S/WW/1

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

FUNGCIDE(2)	O	BENODANI	MEAN		
N SPRING					
0	3.34	3.47	3.40		
NC 1	5.63	5.48	5.55		
NC 2	6.56	6.48	6.52		
NC 3	7.14	6.75	6.94		
MEAN	5.67	5.54	5.60		
FUNGCIDE(2)	O	BENODANI	MEAN		
N SUMMER					
0	5.32	5.42	5.37		
AG 1	6.01	5.67	5.84		
MEAN	5.67	5.54	5.60		
FUNGCIDE(2)	O	BENODANI	MEAN		
FUNGCIDE(1)	O	BENODANI	MEAN		
BN+CA+MA	5.64 5.69	5.39 5.70	5.51 5.69		
MEAN	5.67	5.54	5.60		
EXTRA	NCA1NCD2 7.27	NCA1NCD3 7.69	IBA1NCD2 6.64	IBA1NCD3 7.52	MEAN 7.28

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	N AUTUMN	N SPRING	N SUMMER	FUNGCIDE(1)
SED	0.093	0.132	0.093	0.093
TABLE	FUNGCIDE(2)	EXTRA	N AUTUMN	N SPRING
			N SUMMER	N SUMMER
SED	0.093	0.263	0.132	0.186
TABLE	N AUTUMN	N SPRING	N SUMMER	N AUTUMN
	FUNGCIDE(1)	FUNGCIDE(1)	FUNGCIDE(1)	FUNGCIDE(2)
SED	0.132	0.186	0.132	0.132
TABLE	N SPRING	N SUMMER	FUNGCIDE(1)	N AUTUMN
	FUNGCIDE(2)	FUNGCIDE(2)	FUNGCIDE(2)	N SPRING
SED	0.186	0.132	0.132	0.186

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
WP	10	0.263	4.4

GRAIN MEAN DM% 79.8

PLOT AREA HARVESTED 0.00098