

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1978

[Full Table of Content](#)



---

### 78/R/BE/5 Aphids and Entomophthora - Beans

#### Rothamsted Research

Rothamsted Research (1979) *78/R/BE/5 Aphids and Entomophthora - Beans* ; Yields Of The Field Experiments 1978, pp 412 - 414 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/R/BE/5

SPRING BEANS

APHIDS AND ENTOMOPHTHORA

Object: To study the effects of a range of factors on the incidence of Entomophthora infection of aphids and on the yield of spring beans - Gt. Knott I.

Sponsor: N. Wilding.

Design: One experiment (78/R/BE/5/1) 4 randomised blocks of 3 plots  
One experiment (78/R/BE/5/2) 3 randomised blocks of 8 plots

Whole plot dimensions: 9.22 x 9.14.

Treatments to 78/R/BE/5/1 (all plots irrigated - total 49 mm):

TREATMNT(1)	Biological and chemical control of aphids:
NONE	None
ENT APH	Entomophthora species applied in live infected aphids
PIRIMICA	Insecticide: Pirimicarb at 0.14 kg in 340 l

Treatments to 78/R/BE/5/2 (all plots unirrigated):

TREATMNT(2)	Biological and chemical control of aphids, fungicides to control Entomophthora:
NONE	None
ENT APH	Entomophthora species applied in live infected aphids
CARBARYL	Insecticide: Carbaryl at 1.49 kg
PIRIMICA	Insecticide: Pirimicarb at 0.14 kg
BENOMYL	Fungicide: Benomyl at 0.61 kg
CAPTAFOL	Fungicide: Captafol at 1.74 kg
MANCOZEB	Fungicide: Mancozeb at 1.36 kg
TRIDEMOR	Fungicide: Tridemorph at 0.52 kg

- NOTES: (1) On all treatments except PIRIMICA aphid colonies were artificially established between 31 May, 1978 and 2 June (10-20 colonies were initiated in each 9.14 m length of row).
- (2) All spray treatments were applied in 340 l.
- (3) Sprays were applied on the following dates:- Pirimicarb: 19 June, 1978. Carbaryl: 12, 28 June and 6 July. Mancozeb, captafol, benomyl and tridemorph: 19 and 28 June, 6, 13, 19 and 26 July.
- (4) Irrigation was applied to BE/5/1 at 25 mm on 1 June, 1978 and 8 mm on each of 19 June, 21 and 25 July.

Basal applications: Manures: FYM at 35 t. Weedkillers: Simazine at 1.1 kg in 220 l. Paraquat at 0.6 kg ion in 220 l.

Seed: Minden, sown at 200 kg.

78/R/BE/5

Cultivations, etc.:— Paraquat applied: 8 Nov, 1977. FYM applied, ploughed: 24 Nov. Rotary harrowed: 8 Mar, 1978. Seed sown: 9 Mar. Simazine applied to BE/5/2: 25 Mar, to BE/5/1: 30 Mar. Combine harvested: 25 Sept. Previous crops: Winter barley (BE/5/2), wheat (BE/5/1) 1976, wheat 1977.

NOTE: Weekly assessments of aphid population density and proportion of aphids infected with *Entomophthora* were made from June to August.

78/R/BE/5/1

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT(1)	NONE	ENT APH	PIRIMICA	MEAN
	5.33	5.65	5.22	5.40

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT(1)
-----	
SED	0.448

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	6	0.634	11.7

GRAIN MEAN DM% 81.6

PLOT AREA HARVESTED 0.00244

78/R/BE/5/2

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT(2)	
NONE	4.36
ENT APH	4.89
CARBARYL	4.09
PIRIMICA	5.13
BENOMYL	5.29
CAPTAFOL	4.76
MANCOZEB	5.12
TRIDEMOR	4.40
MEAN	4.75

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT(2)
-----	
SED	0.412

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	14	0.504	10.6

GRAIN MEAN DM% 81.2

PLOT AREA HARVESTED 0.00244