

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1978

[Full Table of Content](#)



---

### 78/R/P/3 Growth Study - Potatoes

#### Rothamsted Research

Rothamsted Research (1979) *78/R/P/3 Growth Study - Potatoes* ; Yields Of The Field Experiments 1978, pp 436 - 437 - DOI: <https://doi.org/10.23637/ERADOC-1-30>

78/R/P/3

POTATOES

GROWTH STUDY

Object: To study the effects of modifying haulm structure by mechanical or chemical means on growth, tuber bulking, radiation interception, photosynthesis rates and yield of potatoes - Long Hoos IV 3.

Sponsor: D.W. Wood.

Design: 2 randomised blocks of 2 plots split into 4.

Whole plot dimensions: 11.6 x 22.0.

Treatments: All combinations of:-

Whole plots

- |             |  |
|-------------|--|
| 1. TBR SIZE | Seed tuber sizes:                      |
| SMALL       | In the weight range 60- 80 g per tuber |
| LARGE       | In the weight range 95-125 g per tuber |

Sub plots

- |             |  |
|-------------|--|
| 2. TREATMNT | Treatments:  |
| NONE        | None   |
| APEX R      | Stem apex removed after tuber initiation   |
| BRANCH R    | Lateral branches removed after tuber initiation  |
| GRTH REG    | Growth regulator, chlorflurenol methyl ester applied as foliar spray at tuber initiation |

NOTE: Chlorflurenol methyl ester applied at 8.4 g in 840 l on 28 June, 1978.

Basal applications: Manures: (0:14:28) at 720 kg. (20:14:14) at 600 kg. 'Nitro-Chalk 25' at 280 kg. Weedkiller: Metribuzin at 0.98 kg in 430 l. Insecticide: Pirimicarb at 0.14 kg. Fungicide: Mancozeb at 1.3 kg in 430 l on the first occasion, applied with the insecticide, at 1.9 kg in 900 l on the second and third occasions.

Seed: Pentland Crown.

Cultivations, etc.: - Autumn PK applied: 4 Oct, 1977. Ploughed: 10 Oct. NPK and additional N applied: 18 Apr, 1978. Heavy spring-tine cultivated: 9 May. Rotary cultivated, ridged, hand planted and split back: 10 May. Rotary ridged: 25 May. Weedkiller applied: 16 June. Insecticide and fungicide applied: 10 July. Fungicide applied: 17 Aug and 28 Aug. Lifted: 9 Oct. Previous crops: Beans 1976, spring wheat 1977.

NOTE: Seed sprout characteristics were assessed at planting. Emergence counts were made daily in early June. Full growth analyses were done at fortnightly intervals between early July and mid-September. Photosynthetic and respiration rates of leaves were assessed. Radiation interception was measured in July.

78/R/P/3

TOTAL TUBERS TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT TBR SIZE	NONE	APEX R	BRANCH R	GRTH REG	MEAN
SMALL	44.3	44.3	45.9	37.3	43.0
LARGE	46.9	43.5	44.7	41.2	44.1
MEAN	45.6	43.9	45.3	39.2	43.5

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT	TBR SIZE* TREATMNT
SED	0.67	0.95

\* WITHIN SAME LEVEL OF TBR SIZE ONLY

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP.SP	6	0.95	2.2

PLOT AREA HARVESTED 0.00181

PERCENTAGE WARE 4.45CM (1.75INCH) RIDDLE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT TBR SIZE	NONE	APEX R	BRANCH R	GRTH REG	MEAN
SMALL	47.0	41.8	31.5	26.8	36.8
LARGE	28.6	22.2	29.7	22.0	25.6
MEAN	37.8	32.0	30.6	24.4	31.2

PLOT AREA HARVESTED 0.00181