

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1979

[Full Table of Content](#)



### 79/W/CS/99 Effects of Breaks on Take-all - Barley, Oats, Beans

#### Rothamsted Research

Rothamsted Research (1980) *79/W/CS/99 Effects of Breaks on Take-all - Barley, Oats, Beans ; Yields Of The Field Experiments 1979*, pp 121 - 122 - DOI: <https://doi.org/10.23637/ERADOC-1-45>

79/W/CS/99

EFFECTS OF BREAKS ON TAKE-ALL

Object: To study factors affecting the incidence of take-all (*Gaeumannomyces graminis*) and effects on yields of barley - Woburn, Butt Furlong.

Sponsor: D. Hornby.

The seventh year, barley, beans, oats.

For previous years see 72/W/CS/99(t) and 73-78/W/CS/99.

Design: 2 blocks of 9 plots.

Whole plot dimensions: 5.33 x 15.2.

Treatments: Crop sequences, soil sterilant and inoculum:-

TREATMNT	1968-71	72	73	74	75	76	77	78	79
B 12	B	B	B	B	B	B	B	B	B
	B	F	B	B	B	B	B	B	F
B 6+S	B	F	BE	B	B	B	B	B	B(S)
B 5	B	B	F	BE	B	B	B	B	B
B 4	B	B	B	F	BE	B	B	B	B
B 3+S+I	B	B	B	B	F	BE	B	B	B(SI)
B 2+I	B	B	B	B	B	F	BE	B	B(I)
	B	B	B	B	B	B	F	BE	O
	B	B	B	B	B	B	B	F	BE

B = Barley, BE = Beans, F = Fallow, O = Oats

S = Soil sterilant: Formalin at 1190 l in 54,200 l

I = Inoculum of take-all applied on colonised autoclaved oats and combine drilled in the ratio of three oats to one barley seed, mixture sown at 310 kg.

Standard applications:

Barley and Oats: Manures: (20:14:14) at 450 kg.

Spring beans: Manures: (0:14:28) at 400 kg. Insecticide: Pirimicarb at 0.14 kg in 220 l.

Seed: Barley: Porthos, dressed with ethirimol, sown at 160 kg.

Oats: Panama, sown at 180 kg.

Spring beans: Minden, sown at 220 kg.

Cultivations, etc.:-

All plots: Ploughed: 17 Nov, 1978. Spring-tine cultivated with crumbler attached: 14 Apr, 1979.

Fallow: Spring-tine cultivated with crumbler attached: 8 May, 1979. Rotary cultivated: 8 June.

Barley: Formalin applied: 18 Apr, 1979. NPK applied, spring-tine cultivated with crumbler attached, seed sown and seed with inoculum combine drilled: 8 May. Combine harvested: 28 Aug.

Oats: NPK applied, spring-tine cultivated, seed sown: 8 May. Harvested by hand: 4 Oct.

Spring beans: Spring-tine cultivated with crumbler attached, PK applied, seed sown: 8 May. Insecticide applied: 11 July. Combine harvested: 10 Sept.

79/W/CS/99

NOTES: (1) Soil samples were taken before treatments were applied for estimates of *Heterodera avenae* cysts and eggs and incidence of microflora.  
(2) Plant samples were taken in July for incidence of 'take-all'.

BARLEY

GRAIN TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

TREATMNT	B 12	B 6+S	B 5	B 4	B 3+S+I	B 2+I	MEAN
	1.97	2.36	2.48	2.34	1.24	1.56	1.99

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	TREATMNT
-----	-----
SED	0.183

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	5	0.183	9.2

GRAIN MEAN DM% 82.3

PLOT AREA HARVESTED 0.00434